

ORBITALLY BUT NOT ASYMPTOTICALLY STABLE GROUND STATES FOR THE DISCRETE NLS

SCIPIO CUCCAGNA

ABSTRACT. We consider examples of discrete nonlinear Schrödinger equations in $\mathbb{Z}$ admitting ground states which are orbitally but not asymptotically stable in $\ell^2(\mathbb{Z})$. The ground states contain internal modes which decouple from the continuous modes. The absence of leaking of energy from discrete to continuous modes leads to an almost conservation and perpetual oscillation of the discrete modes. This is quite different from what is known for nonlinear Schrödinger equations in $\mathbb{R}^d$. We do not investigate connections with work on quasi periodic solutions as in [JA,BV].

§1 INTRODUCTION

We consider the discrete Laplacian Δ in $\mathbb{Z}$ defined by

$$(\Delta u)(n) = u(n+1) + u(n-1) - 2u(n).$$

In $\ell^2(\mathbb{Z})$ we have for the spectrum $\sigma(-\Delta) = [0, 4]$. Let for $\langle n \rangle = \sqrt{1+n^2}$

$$\begin{aligned} \ell^{p,\sigma}(\mathbb{Z}) &= \{u = \{u_n\} : \|u\|_{\ell^{p,\sigma}}^p = \sum_{n \in \mathbb{Z}} \langle n \rangle^{p\sigma} |u(n)|^p < \infty\} \text{ for } p \in [1, \infty) \\ \ell^{\infty,\sigma}(\mathbb{Z}) &= \{u = \{u(n)\} : \|u\|_{\ell^{\infty,\sigma}} = \sup_{n \in \mathbb{Z}} \langle n \rangle^\sigma |u(n)| < \infty\}. \end{aligned}$$

We set $\ell^{p,\sigma} = \ell^{p,\sigma}(\mathbb{Z})$ and $\ell^p = \ell^{p,0}$. We consider a potential $q = \{q(n), n \in \mathbb{Z}\}$ with $q(n) \in \mathbb{R}$ for all n . We consider the discrete Schrödinger operator H

$$(1.1) \quad (Hu)(n) = -(\Delta u)(n) + q(n)u(n).$$

We assume:

- (H1) $|q(n)| \leq Ce^{-|n|}$.
- (H2) The points 0 and 4 are not resonances of H .
- (H3) $\sigma_d(H)$ consists of exactly two eigenvalues $-E_0 < 0$ and $E_1 > 4$.

Here $\lambda = 0$ (resp. 4) is a resonance if there is a nontrivial $u \in \ell^\infty$ with $Hu = \lambda u$. We will see in Appendix A that there are operators satisfying (H1)–(H3). By Lemma

Typeset by $\mathcal{A}\mathcal{M}\mathcal{S}$ -TEX

5.3 [CT], $\dim \ker(H + E_0) = 1$ in ℓ^2 . We denote by $\varphi_0(n)$ a generator of $\ker(H + E_0)$ normalized so that $\|\varphi_0\|_{\ell^2} = 1$. Similarly $\dim \ker(H - E_1) = 1$ in ℓ^2 . We denote by $\varphi_1(n)$ a generator of $\ker(H - E_1)$ normalized so that $\|\varphi_1\|_{\ell^2} = 1$. We pick $\varphi_0(n) > 0$ and $\varphi_1(n) \in \mathbb{R}$ for all n . Consider now the discrete nonlinear Schrödinger equation (DNLS)

$$(1.2) \quad i\partial_t u(t, n) - (Hu)(t, n) + |u(t, n)|^6 u(t, n) = 0.$$

We consider a family of ground states solutions $e^{i\omega t} \phi_\omega$ of (1.2), or equivalently of

$$(1.3) \quad (Hu)(n) - |u(n)|^6 u(n) = -\omega u(n).$$

Specifically we have, see Appendix A [CT]:

Lemma 1.1. *Assume (H1)–(H3). There is a family $\omega \rightarrow \phi_\omega$ of standing waves solving (1.3) with the following properties. For any $\sigma \geq 0$ there is an $\eta > 0$ such that $\omega \rightarrow \phi_\omega$ belongs to $C^\omega([E_0, E_0 + \eta], \ell^{2,\sigma}) \cap C^0([E_0, E_0 + \eta], \ell^{2,\sigma})$. We have $\phi_\omega(n) \in \mathbb{R}$ for any n and there are fixed $a > 0$ and $C > 0$ such that $|\phi_\omega(n)| \leq Ce^{-a|n|}$. As $\omega \searrow E_0$ we have in $C^\infty([E_0, E_0 + \eta], \ell^{2,\sigma}) \cap C^0([E_0, E_0 + \eta], \ell^{2,\sigma})$ the expansion*

$$\phi_\omega = (\omega - E_0)^{\frac{1}{6}} \|\varphi_0\|_{\ell^8}^{-\frac{4}{3}} (\varphi_0 + O(\omega - E_0)).$$

Under our hypotheses, the ground states $e^{i\omega t} \phi_\omega$ are by well known arguments orbitally stable in ℓ^2 . A natural question is whether they are also asymptotically stable. Here we will define asymptotic stability as follows:

Definition 1.2. *We say that an orbitally stable ground state $e^{i\omega_0 t} \phi_{\omega_0}$ of (1.2) is asymptotically stable if there are a $\sigma > 0$ and an $\epsilon_0 > 0$ such that for any $u_0 \in \ell^{1,\sigma}$ with $\|u_0 - \phi_{\omega_0}\|_{\ell^{1,\sigma}} \leq \epsilon_0$ there are a ω_+ and a real valued function $\theta(t)$ with*

$$\lim_{t \rightarrow +\infty} \|u(t) - e^{i\theta(t)} \phi_{\omega_+}\|_{\ell^{\infty, -\sigma}} = 0.$$

Our stating point is the following theorem, proved in [CT], for a weaker result see also [KPS]:

Theorem 1.3. *Suppose that H satisfies (H3), has just one single eigenvalue and $q \in \ell^{1,1}$. Then for any $\omega_0 \in]E_0, E_0 + \eta[$ there exist an $\epsilon_0 > 0$ and a $C > 0$ such that if we pick $u_0 \in \ell^2$ with $\|u_0 - \phi_{\omega_0}\|_{\ell^2} < \epsilon < \epsilon_0$, then there exist $\omega_+ \in (E_0, E_0 + \eta_0)$, $\theta \in C^1(\mathbb{R})$ and $u_+ \in \ell^2$ with $|\omega_+ - \omega_0| + \|u_+\|_{\ell^2} \leq C\epsilon$ such that if $u(t, n)$ is the corresponding solution of (1.2) with $u(0, n) = u_0(n)$, then*

$$\lim_{t \rightarrow \infty} \|u(t) - e^{i\theta(t)} \phi_{\omega_+} - e^{it\Delta} u_+\|_{\ell^2} = 0.$$

Theorem 1.3 implies asymptotic stability of the standing waves $e^{i\omega t} \phi_\omega$ in the sense of Definition 1.2. Theorem 1.3 holds because H is rather special. In fact, consistently with known results in the literature [BV, JA] we show what follows:

Theorem 1.4. Consider in Lemma 1.1 $\sigma > 0$ large and $\eta > 0$ small. Assume (H1)–(H3).

(1) Any $\omega_0 \in]E_0, E_0 + \eta[$ is not asymptotically stable. More precisely for any $\sigma > 0$ there is a sequence with $\|u_n(0) - \phi_{\omega_0}\|_{\ell^{2,\sigma}} \rightarrow 0$ such that for any n

$$\inf_{\omega, \gamma, t} \|u_n(t) - e^{i\gamma} \phi_{\omega}\|_{\ell^{2,-\sigma}} > 0.$$

(2) For $\epsilon > 0$ small enough there is a fixed $A_0(\omega_0)$ such that for $\|u_0 - \phi_{\omega_0}\|_{\ell^2} \leq \epsilon$ we can write for all $t \in \mathbb{R}$ and for a $\theta \in C^1(\mathbb{R}, \mathbb{R})$

$$(1.4) \quad u(t, n) = e^{i\theta(t)}(\phi_{\omega(t)}(n) + r(t, n)), \quad \theta(t) = \int_0^t \omega(t') dt' + \gamma(t),$$

$$(1.5) \quad \text{with } |\omega_0 - \omega(t)| + \|r(t)\|_{\ell^2} \leq A_0(\omega_0)\epsilon$$

$$(1.6) \quad \text{and } \langle \Re r(t), \phi_{\omega(t)} \rangle = \langle \Im r(t), \partial_{\omega} \phi_{\omega(t)} \rangle = 0.$$

(3) We have a representation for all times t

$$(1.7) \quad u(t) = e^{i\theta(t)} (\phi_{\omega(t)} + z\xi_1(\omega(t)) + \bar{z}\xi_2(\omega(t))) + A(\omega(t), z(t)) + h(t)$$

with for a fixed $C = C(\omega_0)$

$$(1.8) \quad \|A(\omega(t), z(t))\|_{\ell^{2,2}} \leq C|z(t)|^2$$

$$(1.9) \quad \lim_{t \rightarrow \infty} \|h(t) - e^{i\Delta t} h_+\|_{\ell^2} = 0 \text{ for a } h_+ \in \ell^2 \text{ with } \|h_+\|_{\ell^2} \leq C\epsilon$$

$$(1.10) \quad \|h\|_{L_t^2 \ell^{2,-2}} \leq C\epsilon$$

$$(1.11) \quad \|\xi_1(\omega) - \varphi_1\|_{\ell^1} + \|\xi_2(\omega)\|_{\ell^1} < C(\omega - E_0).$$

There is another variable $z = \zeta + \alpha(\omega, \zeta) + \mathcal{A}(\omega, \zeta, h)$ with $|\alpha(\omega, \zeta)| \leq C|\zeta|^2$ and $|\mathcal{A}(\omega, \zeta, h)| \leq C|\zeta|\|h(t)\|_{\ell^{2,-2}}$ such that

$$(1.12) \quad i\dot{\zeta} - \lambda(\omega)\zeta = d(\omega, |\zeta|^2)\zeta + \mathcal{B}(\omega, \zeta, h)$$

with $d(\omega, |\zeta|^2)$ real valued and $|\mathcal{B}(\omega, \zeta, h)| \leq C\|h\|_{\ell^{2,-2}}^2$. There is a change of variables $\omega = \varpi + \alpha(\varpi, \zeta) + \mathcal{C}(\varpi, \zeta, h)$ with $|\alpha(\varpi, \zeta)| \leq C|\zeta|^2$ and $\|C(\varpi, \zeta, h)\|_{\ell^{2,2}} \leq C|\zeta|\|h\|_{\ell^{2,-2}}$ such that

$$(1.13) \quad i\dot{\varpi} = D(\varpi, \zeta, h)$$

with $|D(\varpi, \zeta, h)| \leq C\|h\|_{\ell^2, -2}^2$. In particular there is a ϖ_+ with $|\omega_0 - \varpi_+| < C\epsilon^2$ such that

$$(1.14) \quad \lim_{t \rightarrow \infty} (\zeta(t), \varpi(t)) = (0, \varpi_+).$$

For Strichartz estimates for $h(t)$ see §5. Claim (1) is not surprising in view of earlier work on quasiperiodic breathers of the DNLS in the anticontinuous limit, [BV,JA]. Claim (2) is just the standard statement of orbital stability. What to our knowledge is new is the detailed analysis of the long time dynamics near the ground states. We do not push our analysis enough to prove the existence of a manifold of quasiperiodic breathers and to prove its asymptotic stability. Yet maybe Claim (3) is suggestive of such a situation.

Our viewpoint is the same used for continuous NLS in $\mathbb{R}^d$ already used in [CT], see references therein. The result is in some sense opposite, but for reasons consistent with the results for continuous NLS or in [CT]. The difference between Theorems 1.3 and 1.4 is due to differences between the discrete spectra $\sigma_d(H)$. The case of Theorem 1.3 is simpler because $\sigma_d(H)$ consists of just one eigenvalue and as a consequence $\sigma_d(\mathcal{H}_\omega) = \{0\}$ with $\mathcal{H}_\omega$ the linearization in (2.3). However, in analogy to the continuous case, one could prove Theorem 1.3 even in cases when $\sigma_d(H)$ is formed by an arbitrary number of eigenvalues and so $\sigma_d(\mathcal{H}_\omega) \supsetneq \{0\}$. To explain why for H satisfying (H3) Theorem 1.3 fails, we need to discuss the nonlinear coupling between discrete and continuous modes. (1.2) can be rewritten decomposing canonically $u(t)$ into the spectral components associated to the spectral decomposition of $\mathcal{H}_{\omega(t)}$. In particular $u(t)$ is expressed as a ground state plus a reminder R , see §2. Asymptotic stability of ground states corresponds to the fact that $R(t)$ disperses. R is decomposed in various components associated to the spectrum of $\mathcal{H}_\omega$. Thanks to the dispersion theory in [SK, KKK, PS, CT] the continuous spectrum can be thought as stable spectrum. The discrete spectrum $\sigma_d(\mathcal{H}_\omega)$ corresponds to central directions. To see whether or not there is asymptotic stability, we need to study the behavior of the discrete components of R . Notice that in the absence of nonlinear coupling, the discrete components of R would describe periodic motions. In the case of the NLS in $\mathbb{R}^d$, for any $\lambda(\omega) \in \sigma_d(\mathcal{H}_\omega) \setminus \{0\}$ there is a fixed integer n such that $n\lambda(\omega) \in \sigma_e(\mathcal{H}_\omega)$, with σ_e the continuous spectrum. In other words, there is resonance between continuous spectrum and nonzero eigenvalues of $\mathcal{H}_\omega$. This fact seems to be responsible for the asymptotic stability for NLS in $\mathbb{R}^d$, and of orbital instability in the case of standing waves with nodes which are linearly stable, see [C]. Using bifurcation theory as in Lemma 1.1 it is easy to find also examples of DNLS in $\mathbb{Z}^d$ such that there is resonance between continuous spectrum and nonzero eigenvalues of $\mathcal{H}_\omega$. In these cases one expects that energy leaks slowly from the discrete modes to the continuous modes because of resonances, and that continuous modes disperse. In this paper hypothesis (H3) assures that $\sigma_d(\mathcal{H}_\omega) \setminus \{0\} = \{\pm\lambda(\omega)\}$ and that $n\lambda(\omega) \notin \sigma_e(\mathcal{H}_\omega)$ for all integers n . This can be easily manufactured because

$\sigma(-\Delta)$ is a bounded set. By the absence of resonances, we show that discrete and continuous modes decouple. The discrete modes persist in their oscillatory motion. The continuous modes consist of a small component confined in a bounded region of space and of a part which scatters as solutions of the linear constant coefficients Schrödinger equation. To prove this decoupling we need to perform a Siegel normal forms argument. It is likely that the techniques in [BV] can be used to prove the existence of a manifold of quasiperiodic solutions, and that our arguments show that this manifold is a local attractor.

We end with some notation. Given an operator A we set $R_A(z) = (A - z)^{-1}$ its resolvent. We will denote by $\mathcal{S}(\mathbb{Z})$ the set of functions $f(n)$ rapidly decreasing as $|n| \nearrow \infty$. We will denote by $\mathcal{S}(\mathbb{R} \times \mathbb{Z})$ the set of functions $f(t, n)$ rapidly decreasing as (t, n) diverges along with all the derivatives $\partial_t^a f(t, n)$ for $a \in \mathbb{N}$. We will denote by $C_0(\mathbb{Z})$ the set of functions $f(n)$ such that $f(n) = 0$ for n near ∞ . Given two Banach spaces X and Y , $B(X, Y)$ will be the space of bounded linear operators defined in X with values in Y . Set $\text{diag}(a, b)$ for the diagonal 2×2 matrix with (a, b) on the diagonal. Given a matrix A we say that it is real, if it has real valued entries. We denote by ${}^t A$ its transpose.

§2 LINEARIZATION, MODULATION AND SET UP

In Appendix B [CT] it is proved:

Lemma 2.1 (Global well posedness). *The DNLS (1.2) is globally well posed, in the sense that any initial value problem $u(0, n) = u_0(n)$ with $u_0 \in \ell^2$ admits exactly one solution $u(t) \in C^\infty(\mathbb{R}, \ell^2)$. The correspondence $u_0 \rightarrow u(t)$ defines a continuous map $\ell^2 \rightarrow C^\infty([T_1, T_2], \ell^2)$ for any bounded interval $[T_1, T_2]$.*

By the implicit function theorem we get:

Lemma 2.2 (Coordinates near standing waves). *Fix ω_0 close to E_0 . Then there are an $\epsilon_0 > 0$ and a $C_0 > 0$ such that any $u_0 \in \ell^2$ with $\|u_0 - \phi_{\omega_0}\|_{\ell^2} \leq \epsilon \leq \epsilon_0$ can be written in a unique way in the form $u_0 = e^{i\gamma(0)}(\phi_{\omega(0)} + r(0))$ with $|\omega_0 - \omega(0)| + |\gamma(0)| + \|r(0)\|_{\ell^2} \leq C_0\epsilon$ and with $\langle \Re r(0), \phi_{\omega(0)} \rangle = \langle \Im r(0), \partial_\omega \phi_{\omega(0)} \rangle = 0$. The correspondence $u_0 \rightarrow (\gamma(0), \omega(0), r(0))$ is a smooth diffeomorphism.*

By standard arguments one has:

Lemma 2.3 (Orbital Stability). *For $\epsilon_0 > 0$ small enough in Lemma 2.2 there is a fixed $A_0(\omega_0)$ such that for $\|u_0 - \phi_{\omega_0}\|_{\ell^2} \leq \epsilon \leq \epsilon_0$ we can express $u(t)$ as (1.4)–(1.6) for all $t \in \mathbb{R}$.*

Consider the initial datum $u_0(n)$. We consider for all t the decomposition (2.1). When we plough the ansatz (2.1) in (1.2) we obtain

$$(2.1) \quad \begin{aligned} i\partial_t r(t, n) &= (Hr)(t, n) + \omega(t)r(t, n) - 4\phi_{\omega(t)}^6(n)r(t, n) - 3\phi_{\omega(t)}^6(n)\bar{r}(t, n) \\ &+ \dot{\gamma}(t)\phi_{\omega(t)}(n) - i\dot{\omega}(t)\partial_\omega\phi_{\omega(t)}(n) + \dot{\gamma}(t)r(t, n) + N(r(t, n)) \end{aligned}$$

for $N(r(t, n)) = O(r^2(t, n))$. We set ${}^tR = (r, \bar{r})$, ${}^t\Phi = (\phi_\omega, \phi_\omega)$ and we rewrite the above equation as

$$(2.2) \quad iR_t = \mathcal{H}_\omega R + \sigma_3 \dot{\gamma} R + \sigma_3 \dot{\gamma} \Phi - i\dot{\omega} \partial_\omega \Phi + \mathcal{N}(R)$$

where for $\sigma_1 = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$, $\sigma_2 = \begin{bmatrix} 0 & i \\ -i & 0 \end{bmatrix}$ and $\sigma_3 = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}$ we have for $\mathcal{H}_0(\omega) = \sigma_3 [H + \omega]$

$$(2.3) \quad \mathcal{H}_\omega = \mathcal{H}_0(\omega) + V_\omega \text{ with } (V_\omega R)(n) = [-4\sigma_3 + 3i\sigma_2] \phi_\omega^6(n) R(n).$$

We have the following spectral information on $\mathcal{H}_\omega$:

Lemma 2.4. *Pick $\eta > 0$ in Lemma 1.1 very small. Then for any $E_0 < \omega < E_0 + \eta$ we have the following facts:*

(1) *For J the operator defined by $J^t(u_1, u_2) := {}^t(\bar{u}_1, \bar{u}_2)$ we have*

$$\sigma_1 \mathcal{H}_\omega = -\mathcal{H}_\omega \sigma_1, [J, \mathcal{H}_\omega] = 0, \sigma_3 \mathcal{H}_\omega = \mathcal{H}_\omega^* \sigma_3.$$

(2) *The spectrum $\sigma(\mathcal{H}_\omega)$ is symmetric with respect to the coordinate axes.*

(3) *The essential spectrum is $\sigma_e(\mathcal{H}_\omega) = \sigma_e(\mathcal{H}_0(\omega)) = (-4 - \omega, -\omega) \cup (\omega, 4 + \omega)$.*

(4) *For the generalized kernel we have $N_g(\mathcal{H}_\omega) = \text{span}\{\sigma_3 \Phi_\omega, \partial_\omega \Phi_\omega\}$.*

(5) *For the discrete spectrum we have $\sigma_d(\mathcal{H}_\omega) = \{0, \lambda_1(\omega), -\lambda_1(\omega)\}$ with, for a fixed C independent from ω , $|\lambda(\omega) - E_1 - \omega| < C(\omega - E_0)$. We have $\dim \ker(\mathcal{H}_\omega \pm \lambda(\omega)) = 1$ and $N_g(\mathcal{H}_\omega \pm \lambda(\omega)) = \ker(\mathcal{H}_\omega \pm \lambda(\omega))$.*

Proof. (1) follows from the definition of $\mathcal{H}_\omega$ and the fact that $\phi_\omega(n) \in \mathbb{R}$. (2) follows from (1). (3) follows from Weil's essential spectrum theorem, Theorem XIII.14 [RS]. $N_g(\mathcal{H}_\omega) \supseteq \text{span}\{\sigma_3 \Phi_\omega, \partial_\omega \Phi_\omega\}$ follows from computation. The equality is a consequence of elementary perturbation theory and the fact that $\mathcal{H}_0(\omega)$ has simple eigenvalues in $\pm(\omega - E_0)$ and no other eigenvalues near 0. The existence of $\lambda(\omega)$ satisfying the above estimates follows from perturbation theory and the fact that $\mathcal{H}_0(\omega)$ has two simple eigenvalues $\pm(E_1 + \omega)$. The fact that $\dim \ker(\mathcal{H}_\omega \pm \lambda(\omega)) = 1$ and $N_g(\mathcal{H}_\omega \pm \lambda(\omega)) = \ker(\mathcal{H}_\omega \pm \lambda(\omega))$ follow from elementary perturbation theory. Finally, for η small enough, the hypotheses (H2) and (H3) imply that $\mathcal{H}_\omega$ does not have other eigenvalues. This ends the proof of Lemma 2.4.

We set $\ell_d^2(\mathcal{H}_\omega) = N_g(\mathcal{H}_\omega) \oplus \oplus_\pm N(\mathcal{H}_\omega \mp \lambda(\omega))$ and $\ell_c^2(\mathcal{H}_\omega) = \{\ell_c^2(\mathcal{H}_\omega^*)\}^\perp$. We have the $\mathcal{H}_\omega$ invariant Jordan block decomposition

$$(2.4) \quad \ell^2 = N_g(\mathcal{H}_\omega) \oplus (\oplus_\pm N(\mathcal{H}_\omega \mp \lambda(\omega))) \oplus \ell_c^2(\mathcal{H}_\omega) = N_g(\mathcal{H}_\omega) \oplus N_g^\perp(\mathcal{H}_\omega^*).$$

We have:

Lemma 2.5. $V_\omega \in \ell^{p,\sigma}$, for any $p \in [1, \infty]$ and $\sigma \in \mathbb{R}$, and $\lambda(\omega)$ depend analytically on $\omega \in]E_0, E_0 + \eta[$. It is possible to choose generators $\xi(\omega) \in \ker(\mathcal{H}_\omega - \lambda(\omega))$ so that $\xi(\omega) \in \ell^{p,\sigma}$, for any $p \in [1, \infty]$ and $\sigma \in \mathbb{R}$, depend analytically on $\omega \in]E_0, E_0 + \eta[$ and we have the normalization $\langle \xi(\omega), \sigma_3 \xi(\omega) \rangle = 1$. Additionally we can choose ${}^t\xi(\omega) = (\xi_1(\omega), \xi_2(\omega))$ so that for any $\tau \in \mathbb{R}$ there is C_τ such that $\|\xi_1 - \varphi_1\|_{\ell^{1,\tau}} < C_\tau(\omega - E_0)$ and $\|\xi_2\|_{\ell^{1,\tau}} < C_\tau(\omega - E_0)$, where φ_1 is a normalized generator of $\ker(H - E_1)$.

Proof. The fact that $V_\omega \in \ell^{p,\sigma}$ depends analytically on ω for $|\omega - \omega_0| \leq \alpha_0$ for some $\alpha_0 > 0$ follows by formula (2.3) and Lemma 1.1. Then $\omega \rightarrow \mathcal{H}_\omega$ is analytic from $|\omega - \omega_0| \leq \alpha_0$ into $B(\ell^{p,\sigma}, \ell^{p,\sigma})$ for all $p \geq 1$ and $\sigma \in \mathbb{R}$. In particular, $\mathcal{H}_\omega$ is an analytic family of operators in the sense of Kato, p. 14 [RS] vol.IV (that is for any fixed ω_0 and $z \notin \sigma(\mathcal{H}_{\omega_0})$, then $R_{\mathcal{H}_\omega}(z)$ is analytic in ω for ω sufficiently close to ω_0). Then by the Kato-Rellich theorem, Theorem XII.8 [RS], $\lambda(\omega)$ depends analytically on ω . Furthermore, we have the projection operator

$$P_{\ker(\mathcal{H}_\omega - \lambda(\omega))} = \frac{i}{2\pi} \int_{|z - E_1 - \omega_0| = a_0} R_{\mathcal{H}_\omega}(z) dz, \quad a_0 = \frac{E_1 - 4}{2}.$$

So $P_{\ker(\mathcal{H}_\omega - \lambda(\omega))}$ depends analytically on ω .

$$\begin{aligned} P_{\ker(\mathcal{H}_\omega - \lambda(\omega))} &= P_{\ker(\mathcal{H}_0 - E_1 - \omega)} + T(\omega) = \\ &= \frac{i}{2\pi} \int_{|z - E_1 - \omega_0| = a_0} (R_{\mathcal{H}_0}(z) - R_{\mathcal{H}_0}(z) V_\omega (1 + R_{\mathcal{H}_0}(z) V_\omega)^{-1} R_{\mathcal{H}_0}(z)) dz, \end{aligned}$$

with, for any $p, q \in [1, \infty]$ and $\tau, \sigma \in \mathbb{R}$, $\|T(\omega)\|_{B(\ell^{p,\tau}, \ell^{p,\sigma})} \lesssim (\omega - E_0)$ with a fixed constant. This implies also the information for ξ and concludes Lemma 2.5.

The conditions $\langle \Re r(t), \phi_{\omega(t)} \rangle = \langle \Im r(t), \partial_\omega \phi_{\omega(t)} \rangle = 0$ are the same of $\langle R(t), \Phi_{\omega(t)} \rangle = \langle R(t), \sigma_3 \partial_\omega \Phi_{\omega(t)} \rangle = 0$, that is $R(t) \in N_g^\perp(\mathcal{H}_{\omega(t)}^*)$. This in particular implies that, in correspondence to the spectral decomposition (2.5) below, we have a decomposition

$$(2.5) \quad R(t) = (z\xi + \bar{z}\sigma_1\xi) + f(t) \in \left[\sum_{\pm} \ker(\mathcal{H}_{\omega(t)} \mp \lambda(\omega(t))) \right] \oplus L_c^2(\mathcal{H}_{\omega(t)}).$$

$R(t) \in N_g^\perp(\mathcal{H}_{\omega(t)}^*)$ implies, for $P_{N_g(\mathcal{H}_\omega)}$ the projection on $N_g(\mathcal{H}_\omega)$ associated to (2.4),

$$(2.6) \quad \begin{aligned} i\dot{\omega} \langle \Phi, \partial_\omega \Phi \rangle &= \langle \sigma_3 \dot{\gamma} R + \mathcal{N}(R) + i\dot{\omega} \partial_\omega P_{N_g(\mathcal{H}_\omega)} R, \Phi \rangle \\ \dot{\gamma} \langle \Phi, \partial_\omega \Phi \rangle &= -\langle \text{same as above}, \sigma_3 \partial_\omega \Phi \rangle. \end{aligned}$$

We have:

Lemma 2.6. *There are two functions $\mu(\omega, r, \bar{r})$ and $\nu(\omega, r, \bar{r})$ defined in the subset of $\ell^2 \times \ell^2$ defined by $|z| + |\bar{z}| + \|f\|_{\ell^2, -\tau} \leq (\omega - E_0)^{\frac{1}{6}} \alpha_0$ and by $|\omega - \omega_0| \leq \beta_0$ for some fixed small $\alpha_0 > 0$ and $0 < \beta_0 \ll (\omega_0 - E_0)$, analytic in $(\omega, r, \bar{r})$, such that for any $\tau \geq 0$ we have $|\nu(\omega, r, \bar{r})| + |\mu(\omega, r, \bar{r})| \leq C(\tau, \omega) \|(r, \bar{r})\|_{\ell^2, -\tau}^2$ with $C(\tau, \omega)$ continuous and with for real ω*

$$(2.7) \quad \begin{aligned} i\dot{\omega} &= i\dot{\omega}(\omega, r, \bar{r}) = \nu(\omega, r, \bar{r}) - \nu(\omega, \bar{r}, r) \text{ with } \overline{\nu(\omega, r, \bar{r})} = \nu(\bar{r}, r) \\ \dot{\gamma} &= \dot{\gamma}(\omega, r, \bar{r}) = \mu(\omega, r, \bar{r}) + \mu(\omega, \bar{r}, r) \text{ with } \overline{\mu(\omega, r, \bar{r})} = \mu(\omega, \bar{r}, r). \end{aligned}$$

Proof. Apply $P_{N_g(\mathcal{H}_\omega)}$ to (2.2) obtaining

$$iP_{N_g(\mathcal{H}_\omega)}R_t - \dot{\gamma}(t)P_{N_g(\mathcal{H}_\omega)}\sigma_3R - \sigma_3\dot{\gamma}(t)\Phi_\omega + i\dot{\omega}(t)\partial_\omega\Phi_\omega + P_{N_g(\mathcal{H}_\omega)}\mathcal{N}(R).$$

Set $q(\omega) = \|\phi_\omega\|_2^2$ and $q'(\omega) = dq(\omega)/d\omega$. Then we have

$$P_{N_g(\mathcal{H}_\omega)} = \sigma_3\Phi_\omega \langle \cdot, \sigma_3\partial_\omega\Phi_\omega \rangle / q'(\omega) + \partial_\omega\Phi_\omega \langle \cdot, \Phi_\omega \rangle / q'(\omega).$$

By $P_{N_g(\mathcal{H}_\omega)}R = 0$, which implies $P_{N_g(\mathcal{H}_\omega)}R_t = -\dot{\omega}\partial_\omega P_{N_g(\mathcal{H}_\omega)}R$, we get

$$\left(q'(\omega) + \begin{bmatrix} -\langle \partial_\omega P_{N_g(\mathcal{H}_\omega)}R, \Phi_\omega \rangle & \langle \sigma_3R, \Phi_\omega \rangle \\ -\langle \partial_\omega P_{N_g(\mathcal{H}_\omega)}R, \sigma_3\partial_\omega\Phi_\omega \rangle & \langle R, \partial_\omega\Phi_\omega \rangle \end{bmatrix} \right) \begin{bmatrix} i\dot{\omega} \\ -\dot{\gamma} \end{bmatrix} = \begin{bmatrix} \langle \mathcal{N}(R), \Phi_\omega \rangle \\ \langle \mathcal{N}(R), \sigma_3\partial_\omega\Phi_\omega \rangle \end{bmatrix}.$$

By an elementary computation we have

$$(2.8) \quad \begin{aligned} \langle \partial_\omega P_{N_g(\mathcal{H}_\omega)}R, \Phi_\omega \rangle &= \langle r + \bar{r}, \partial_\omega\phi_\omega \rangle \\ \langle \partial_\omega P_{N_g(\mathcal{H}_\omega)}R, \sigma_3\partial_\omega\Phi_\omega \rangle &= \langle r - \bar{r}, \partial_\omega^2\phi_\omega \rangle \end{aligned}$$

and so ${}^t(i\dot{\omega}, -\dot{\gamma}) =$

$$(2.9) \quad \left(q'(\omega) + \begin{bmatrix} -\langle r + \bar{r}, \partial_\omega\phi_\omega \rangle & \langle r - \bar{r}, \phi_\omega \rangle \\ -\langle r - \bar{r}, \partial_\omega^2\phi_\omega \rangle & \langle r + \bar{r}, \partial_\omega\phi_\omega \rangle \end{bmatrix} \right)^{-1} \begin{bmatrix} \langle N(r, \bar{r}) - N(\bar{r}, r), \phi_\omega \rangle \\ \langle N(r, \bar{r}) + N(\bar{r}, r), \partial_\omega\phi_\omega \rangle \end{bmatrix}.$$

Notice that the right factor in (2.9) is a polynomial in $(r, \bar{r})$ while the left factor is analytic in the functions in (2.8) and $\langle r - \bar{r}, \phi_\omega \rangle$. This completes Lemma 2.6.

We plug now decomposition (2.6) in system (2.3) and we obtain

$$(2.10) \quad \begin{aligned} i\dot{z} - \lambda(\omega)z &= \langle \dot{\gamma}\sigma_3R + \mathcal{N}(R) - \\ &\quad - iz\dot{\omega}\partial_\omega\xi - i\bar{z}\dot{\omega}\sigma_1\partial_\omega\xi + i\dot{\omega}\partial_\omega P_{\ker(\mathcal{H}_\omega - \lambda)}R, \sigma_3\xi \rangle \\ if - \mathcal{H}_\omega f &= \dot{\gamma}P_c(\mathcal{H}_\omega)\sigma_3R + P_c(\mathcal{H}_\omega)\mathcal{N}(R) + i\dot{\omega}\partial_\omega P_c(\mathcal{H}_\omega)R. \end{aligned}$$

Our first step will consist in §4 in splitting $f = \Phi(\omega, z) + g$ with $g \in \ell_c^2(\mathcal{H}_\omega)$ satisfying an equation of the form (4.1) below and with $\|\Phi(\omega, z)\|_{\ell^{2,2}} \leq C|z|^2$ for fixed $C > 0$. In §5 we will prove that g is asymptotically free. In §6 we will prove that $z(t)$ does not decay to 0. We will also show in §7 that $\omega(t)$ oscillates. We first state some linear dispersive estimates needed later. Proofs of Lemmas 3.1 and 3.3 are in sections 8–10.

§3 SPACETIME ESTIMATES FOR $\mathcal{H}_\omega$

We list a number of linear estimates needed later. The constants $C(\omega)$ and $C(\tau, \omega)$ in this section are upper semicontinuous in ω .

Lemma 3.1. *Under hypotheses (H1-3) there is a constant $C(\omega)$ such that*

$$\|P_c(\mathcal{H}_\omega)e^{it\mathcal{H}_\omega}\|_{B(\ell^p, \ell^{p'})} \leq C(\omega)\langle t \rangle^{-\frac{2}{3}(\frac{1}{p}-\frac{1}{2})} \quad \forall p \in [1, 2] \text{ and for } p' = \frac{p}{p-1}.$$

The proof is in §10. The next estimates needed are Strichartz estimates. Following [CV] for every $1 \leq p, q \leq \infty$ we introduce the Birman-Solomjak spaces

$$\ell^p(\mathbb{Z}, L_t^q[n, n+1]) \equiv \{f \in L_{loc}^q(\mathbb{R}) \text{ s.t. } \{\|f\|_{L^q[n, n+1]}\}_{n \in \mathbb{Z}} \in \ell^p(\mathbb{Z})\},$$

endowed with the norms

$$\begin{aligned} \|f\|_{\ell^p(\mathbb{Z}, L_t^q[n, n+1])}^p &\equiv \sum_{n \in \mathbb{Z}} \|f\|_{L_t^q[n, n+1]}^p \quad \forall \quad 1 \leq p < \infty \text{ and } 1 \leq q \leq \infty \\ \|f\|_{\ell^\infty(\mathbb{Z}, L_t^q[n, n+1])} &\equiv \sup_{n \in \mathbb{Z}} \|f\|_{L^q[n, n+1]}. \end{aligned}$$

We say that a pair of numbers (r, p) is admissible if

$$(3.1) \quad 2/r + 1/p = 1/2 \text{ and } (r, p) \in [4, \infty] \times [2, \infty].$$

Then proceeding as in [CV], by a standard TT^* argument it is possible to prove from Lemma 3.1 the following result:

Lemma 3.2 (Strichartz estimates). *Under the hypotheses of Theorem 1.3 there exists a constant $C = C(\omega)$ upper semicontinuous in ω such that for every admissible pair (r, p) we have:*

$$\|e^{it\mathcal{H}_\omega} P_c(\mathcal{H}_\omega)f\|_{\ell^{\frac{3}{2}r}(\mathbb{Z}, L_t^\infty([n, n+1], \ell^p(\mathbb{Z})))} \leq C(\omega)\|f\|_{\ell^2(\mathbb{Z})}.$$

Moreover, for any two admissible pairs $(r_1, p_1), (r_2, p_2)$ we have the estimate

$$\begin{aligned} &\left\| \int_0^t e^{i(t-s)\mathcal{H}_\omega} P_c(\mathcal{H}_\omega)g(s)ds \right\|_{\ell^{\frac{3}{2}r_1}(\mathbb{Z}, L_t^\infty([n, n+1], \ell^{p_1}(\mathbb{Z})))} \leq \\ &\leq C(\omega)\|g\|_{\ell^{(\frac{3}{2}r_2)'}(\mathbb{Z}, L_t^1([n, n+1], \ell^{p_2'}(\mathbb{Z})))}. \end{aligned}$$

In §9 we prove the following Kato smoothness result:

Lemma 3.3. For $\tau > 1$ there exists $C = C(\tau, \omega)$ such that for all $z \in \mathbb{C} \setminus \sigma_e(\mathcal{H}_\omega)$

$$\|R_{\mathcal{H}_\omega}(z)P_c(\mathcal{H}_\omega)\|_{B(\ell^{2,\tau}, \ell^{2,-\tau})} \leq C.$$

The following limits are well defined for any $\lambda \in [0, 4]$ in $C^0([0, 4], B(\ell^{2,\tau}, \ell^{2,-\tau}))$

$$\lim_{\epsilon \rightarrow 0^+} R_{\mathcal{H}_\omega}(\lambda \pm i\epsilon) = R_{\mathcal{H}_\omega}^\pm(\lambda).$$

For any $u \in \ell^{2,\tau} \cap \ell_c^2(\mathcal{H}_\omega)$ we have

$$\begin{aligned} P_c(\mathcal{H}_\omega)u &= \frac{1}{\sqrt{2\pi}i} \int_{\mathbb{R}} (R_{\mathcal{H}_\omega}^+(\lambda) - R_{\mathcal{H}_\omega}^-(\lambda))u d\lambda \\ &= \frac{1}{\sqrt{2\pi}i} \int_{\sigma_e(\mathcal{H}_\omega)} (R_{\mathcal{H}_\omega}^+(\lambda) - R_{\mathcal{H}_\omega}^-(\lambda))u d\lambda. \end{aligned}$$

The first two statements are proved in Lemma 9.1. The third statement is proved in Lemma 9.4. We list now a number of corollaries.

Lemma 3.4. For $\tau > 3/2 \exists C = C(\tau, \omega)$ s.t.:

(a) for any $f \in \mathcal{S}(\mathbb{Z})$,

$$\|e^{-it\mathcal{H}_\omega}P_c(\mathcal{H}_\omega)f\|_{L_t^2\ell^{2,-\tau}} \leq C\|f\|_{\ell^2};$$

(b) for any $g(t, n) \in \mathcal{S}(\mathbb{R} \times \mathbb{Z})$

$$\left\| \int_{\mathbb{R}} e^{it\mathcal{H}_\omega}P_c(\mathcal{H}_\omega)g(t, \cdot)dt \right\|_{\ell^2} \leq C\|g\|_{L_t^2\ell^{2,\tau}}.$$

The proof is the same of Lemma 3.3 [CT].

Lemma 3.5. For any $\tau > 1 \exists C = C(\tau, \omega)$ such that

$$\left\| \int_0^t e^{-i(t-s)\mathcal{H}_\omega}P_c(\mathcal{H}_\omega)g(s, \cdot)ds \right\|_{L_t^2\ell^{2,-\tau}} \leq C\|g\|_{L_t^2\ell^{2,\tau}}.$$

The proof is the same of Lemma 3.4 [CT].

Lemma 3.6. For every $\tau > 3/2 \exists C = C(\tau, \omega)$ such that

$$\left\| \int_0^t e^{-i(t-s)\mathcal{H}_\omega}P_c(\mathcal{H}_\omega)g(s, \cdot)ds \right\|_{L_t^\infty\ell^2 \cap \ell^6(\mathbb{Z}, L_t^\infty([n, n+1], \ell^\infty))} \leq C\|g\|_{L_t^2\ell^{2,\tau}}.$$

The proof is the same of Lemma 3.5 [CT]. We will now assume Lemmas 3.1 and 3.3 and we will proceed with the proof of our nonlinear result.

§4 DECOUPLING BETWEEN LOCALIZED AND DISPERSIVE RADIATION

Lemma 4.1. *There is a representation $f = \Phi(\omega, z) + g$ such that $\sigma_1 g = \bar{g}$, $\Phi(\omega, z) \in \ell_c^2(\mathcal{H}_\omega)$ is analytic in $(\omega, z, \bar{z})$ with $\|\Phi(\omega, z)\|_{\ell^{2,2}} \leq C|z|^2$ for fixed $C > 0$, and $g \in \ell_c^2(\mathcal{H}_\omega)$ which satisfies, for $\|\widehat{\mathcal{F}}(\omega, z, g)\|_{\ell^{2,2}} \leq C|z| \|g\|_{\ell^{2,-2}}$,*

$$(4.1) \quad i\dot{g} - (\mathcal{H}_\omega + \dot{\gamma}P_c(\mathcal{H}_\omega)\sigma_3)g = \widehat{\mathcal{F}}(\omega, z, g) + O(|g|^7).$$

Proof. We enter the splitting (2.6) in equations (2.7) and (2.10). We set $f_1 = f$ and setting $\dot{\omega} = \dot{\omega}(\omega, R)$ $\dot{\gamma} = \dot{\gamma}(\omega, R)$, i.e. the functions in Lemma 2.6, we write

$$(4.2) \quad \begin{aligned} i\dot{f}_1 - \mathcal{H}_\omega f_1 &= \dot{\gamma}P_c(\mathcal{H}_\omega)\sigma_3 f_1 + A_1(\omega, z) + \mathcal{F}_1(\omega, z, f_1) \\ i\dot{\omega} &= b_1(\omega, z) + \Omega_1(\omega, z, f_1) \\ \dot{\gamma} &= c_1(\omega, z) + \Gamma_1(\omega, z, f_1) \\ i\dot{z} - \lambda(\omega)z &= d_1(\omega, z) + Z_1(\omega, z, f_1). \end{aligned}$$

We will define recursively a sequence of systems

$$(4.3) \quad \begin{aligned} i\dot{f}_\ell - \mathcal{H}_\omega f_\ell &= \dot{\gamma}P_c(\mathcal{H}_\omega)\sigma_3 f_\ell + A_\ell(\omega, z) + \mathcal{F}_\ell(\omega, z, f_\ell) \\ i\dot{\omega} &= b_\ell(\omega, z) + \Omega_\ell(\omega, z, f_\ell) \\ \dot{\gamma} &= c_\ell(\omega, z) + \Gamma_\ell(\omega, z, f_\ell) \\ i\dot{z} - \lambda(\omega)z &= d_\ell(\omega, z) + Z_\ell(\omega, z, f_\ell) \end{aligned}$$

and we will assume a number of **inductive hypotheses** for fixed $\varepsilon_0 > 0$ small enough and C :

- (1) $f_\ell \in \ell_c^2(\mathcal{H}_\omega)$;
- (2) $\mathcal{F}_\ell(\omega, z, f_\ell) = \mathcal{F}(f_\ell) + \widehat{\mathcal{F}}_\ell(\omega, z, f_\ell)$ with $\mathcal{F}(f_\ell) = O(|f_\ell|^7)$ and with $\widehat{\mathcal{F}}_\ell(\omega, z, f_\ell) \in \ell^{2,2}$;
- (3) $b_\ell, c_\ell, d_\ell, \Omega_\ell, \Gamma_\ell$ and Z_ℓ are analytic functions in $(\omega, z, \bar{z}, f_\ell)$ for

$$(4.4) \quad \max\{|\omega - \omega_0|, |z|, \|f_\ell\|_{\ell^{2,-2}}\} \leq e^{-(2-2^{-\ell})}\varepsilon_0;$$

furthermore, their Taylor expansions have real coefficients;

- (4) $(\omega, z, f_\ell) \rightarrow \widehat{\mathcal{F}}_\ell(\omega, z, f_\ell) \in \ell^{2,2}$ is an analytic function in the domain in (4);
- (5) $(\omega, z) \rightarrow A_\ell(\omega, z) \in \ell^{2,2}$ is analytic in $(\omega, z, \bar{z}, f_\ell)$ for $\max\{|\omega - \omega_0|, |z|\} \leq e^{-(2-2^{-\ell})}\varepsilon_0$; if we expand

$$A_\ell(\omega, z) = \sum_{m+n \geq \ell+1} A_{\ell mn}(\omega) z^m \bar{z}^n \text{ and set } \widetilde{A}_\ell(\omega, z) := \sum_{m+n \geq \ell+2} A_{\ell mn}(\omega) z^m \bar{z}^n,$$

- then the $A_{\ell mn}(\omega)$ are real with $\sigma_1 A_{\ell mn}(\omega) = -A_{\ell nm}(\omega)$; $A_\ell(\omega, z) \in \ell_c^2(\mathcal{H}_\omega) \cap \ell^{2,2}$;
(6) The subspace of solutions of (4.3) with ω real, z and $\bar{z}$ complex conjugate of each other and $\sigma_1 f_\ell = \bar{f}_\ell$, is invariant by system (4.3);
(7) the following estimates hold in (4.4)

$$(4.5) \quad \begin{aligned} & \|A_\ell(\omega, z)\|_{\ell^{2,2}} \leq C_A(\ell) e^{(\ell-1)(2-2^{-\ell})} \varepsilon_0^{-\ell+1} |z|^{\ell+1} \\ & \max\{(|\Omega_\ell|, |\Gamma_\ell|, |Z_\ell|, \|\widehat{\mathcal{F}}_\ell\|_{\ell^{2,2}})(\omega, z, f_\ell)\} + \leq C_\Omega(\ell) (|z| + \|f_\ell\|_{\ell^{2,-2}}) \|f_\ell\|_{\ell^{2,-2}} \\ & \max\{|b_\ell(\omega, z)|, |c_\ell(\omega, z)|, |d_\ell(\omega, z)|\} \leq C_b(\ell) |z|^2. \end{aligned}$$

These hypotheses hold for $\ell = 1$, for $C_A(1) = C_\Omega(1) = C_b(1) = c(1)$. We define for $\Phi_{(\ell+1)mn}(\omega) := R_{\mathcal{H}_\omega}((m-n)\lambda(\omega))A_{\ell mn}(\omega)$

$$f_\ell = f_{\ell+1} + \Phi_{\ell+1}(\omega, z), \quad \Phi_{\ell+1}(\omega, z) = \sum_{m+n=\ell+1} \Phi_{(\ell+1)mn}(\omega) z^m \bar{z}^n.$$

We have $\Phi_{\ell+1} \in \ell_c^2(\mathcal{H}_\omega)$. This implies $f_{\ell+1} \in \ell_c^2(\mathcal{H}_\omega)$. We have:

Lemma 4.2. $\Phi_{\ell+1}(\omega, z)$ is analytic in $(\omega, z, \bar{z})$ for $|\omega - \omega_0| \leq e^{-(2-2^{-\ell})}\varepsilon_0$ and $(z, \bar{z}) \in \mathbb{C}^2$ with values in $\ell^{2,2}$. In $|\omega - \omega_0| \leq e^{-(2-2^{-\ell-1})}\varepsilon_0$ and for $\Phi'_{\ell+1} \cdot (b, c) := \partial_\omega \Phi_{\ell+1} b + \partial_z \Phi_{\ell+1} c - \partial_{\bar{z}} \bar{\Phi}_{\ell+1} \bar{c}$ we have for $D_\ell = C_{4.9} \frac{e^{2 \cdot 2^{\ell+2}}}{\ell!}$, see (4.9) for $C_{4.9}$,

$$(4.6) \quad \begin{aligned} & \|\Phi_{\ell+1}(\omega, z)\|_{\ell^{2,2}} \leq C_A(\ell) D_\ell e^{(\ell-1)(2-2^{-\ell})} \varepsilon_0^{-\ell+1} |z|^{\ell+1} \leq C_A(\ell) D_\ell |z|^2 \\ & \|\Phi'_{\ell+1}(\omega, z)\|_{\ell^{2,2}} \leq 4C_A(\ell) D_\ell e^{(\ell-1)(2-2^{-\ell})} \varepsilon_0^{-\ell+1} |z|^\ell \leq 4C_A(\ell) D_\ell |z|. \end{aligned}$$

For ω real $\Phi_{(\ell+1)mn}(\omega)$ is real, $\sigma_1 \Phi_{(\ell+1)mn}(\omega) = \Phi_{(\ell+1)nm}(\omega)$, and so $\sigma_1 f_{\ell+1} = \overline{f_{\ell+1}}$ if $\sigma_1 f_\ell = \bar{f}_\ell$. In particular this yields the inductive hypothesis (6) for $\ell + 1$.

Proof. The last two sentences follow from $\sigma_1 A_{\ell mn}(\omega) = -A_{\ell mn}(\omega)$ for ω real and from $\sigma_1 \mathcal{H}_\omega = -\mathcal{H}_\omega \sigma_1$. Now we turn to the estimates. There are fixed $C_0 > 0$ and $\alpha_0 > 0$ such that for $|\omega - \omega_0| \leq \varepsilon_0$ and $j = 0, 1$

$$(4.7) \quad |\partial_\omega^j [R_{\mathcal{H}_0(\omega)}((m-n)\lambda(\omega), \mu, \nu)]| \leq C_0 e^{-\alpha_0 |\mu - \nu|}.$$

Set $P_c(\omega) = P_c(H_\omega)$ and $\mathcal{H}_\omega^c = P_c(\omega) \mathcal{H}_\omega$. Then

$$(4.8) \quad R_{\mathcal{H}_\omega^c} = R_{\mathcal{H}_0(\omega)} P_c(\omega) - R_{\mathcal{H}_0(\omega)} V_\omega P_c(\omega) R_{\mathcal{H}_0(\omega)} + R_{\mathcal{H}_0(\omega)} V_\omega R_{\mathcal{H}_\omega^c} V_\omega R_{\mathcal{H}_0(\omega)}.$$

By (4.7)–(4.8) for $|\omega - \omega_0| \leq \varepsilon_0$ and for all $(m, n) \in \mathbb{Z}^2$

$$(4.9) \quad \|\partial_\omega^j [R_{\mathcal{H}_\omega^c}((m-n)\lambda(\omega))]\|_{B(\ell^{2,2}, \ell^{2,2})} \leq C_{4.9} \text{ for } j = 0, 1.$$

(4.9) implies for $\max\{|\omega - \omega_0|, |z|\} \leq e^{-(2-2^{-\ell-1})}\varepsilon_0$

$$\begin{aligned}
(4.10) \quad & \|\Phi_{\ell+1}\|_{\ell^2,2} \leq C_{4.9} \sum_{m+n=\ell+1} \frac{1}{m!n!} \|\partial_z^m \partial_{\bar{z}}^n A_\ell(\omega, 0)\|_{\ell^2,2} |z|^{\ell+1} \\
& \leq \frac{C_{4.9} 2^{\ell+1} |z|^{\ell+1}}{(\ell+1)! (\varepsilon_0 \exp(2^{-\ell} - 2))^{\ell+1}} \sup_{|z| \leq e^{-(2-2^{-\ell})}\varepsilon_0} \|A_\ell(\omega, z)\|_{\ell^2,2} \\
& \leq C_{4.9} C_A(\ell) \frac{2^{\ell+1} |z|^{\ell+1}}{(\ell+1)! (\varepsilon_0 \exp(2^{-\ell} - 2))^{\ell-1}}.
\end{aligned}$$

Differentiating we obtain $\partial_\omega \Phi_{\ell+1} = \Phi_{\ell+1}^{(1)} + \Phi_{\ell+1}^{(2)}$ with

$$\begin{aligned}
\Phi_{\ell+1}^{(1)}(\omega, z) &= \sum_{m+n=\ell+1} \partial_\omega [R_{\mathcal{H}_\omega^c}((m-n)\lambda(\omega))] A_{\ell mn}(\omega) z^m \bar{z}^n \\
\Phi_{\ell+1}^{(2)}(\omega, z) &= \sum_{m+n=\ell+1} R_{\mathcal{H}_\omega^c}((m-n)\lambda(\omega)) \partial_\omega A_{\ell mn}(\omega) z^m \bar{z}^n.
\end{aligned}$$

We have $\|\Phi_{\ell+1}^{(1)}(\omega, z)\|_{\ell^2,2} \leq C_{4.9} C_A(\ell) \frac{2^{\ell+1} |z|^{\ell+1}}{(\ell+1)! (\varepsilon_0 \exp(2^{-\ell} - 2))^{\ell-1}}$ as for (4.10). By

$$(4.11) \quad \varepsilon_0 \exp(2^{-\ell} - 2) - \varepsilon_0 \exp(2^{-\ell-1} - 2) \geq \varepsilon_0 e^{-2} 2^{-\ell-1}$$

and by the Cauchy integral formula, we have

$$\begin{aligned}
\|\Phi_{\ell+1}^{(2)}(\omega, z)\|_{\ell^2,2} &\leq C_{4.9} \sum_{m+n=\ell+1} \frac{1}{m!n!} \|\partial_z^m \partial_{\bar{z}}^n \partial_\omega A_\ell(\omega, 0)\|_{\ell^2,2} |z|^{\ell+1} \\
&\leq C_{4.9} C_A(\ell) \frac{2^{2\ell+2} e^2 |z|^{\ell+1}}{(\ell+1)! \varepsilon_0 (\varepsilon_0 \exp(2^{-\ell} - 2))^{\ell-1}}.
\end{aligned}$$

So we conclude

$$(4.12) \quad \|\partial_\omega \Phi_{\ell+1}(\omega, z)\|_{\ell^2,2} \leq 2C_{4.9} C_A(\ell) \frac{2^{2\ell+2} e^2 |z|^{\ell+1}}{(\ell+1)! \varepsilon_0 (\varepsilon_0 \exp(2^{-\ell} - 2))^{\ell-1}}.$$

We have

$$\begin{aligned}
(4.13) \quad & \|\partial_z \Phi_{\ell+1}\|_{\ell^2,2} \leq C_{4.9} \sum_{m+n=\ell+1} \frac{1}{(m-1)!n!} \|\partial_z^m \partial_{\bar{z}}^n A_\ell(\omega, 0)\|_{\ell^2,2} |z|^\ell \\
& \leq C_{4.9} C_A(\ell) \frac{2^{\ell+1} |z|^\ell}{\ell! (\varepsilon_0 \exp(2^{-\ell} - 2))^{\ell-1}}.
\end{aligned}$$

This concludes the proof of Lemma 4.2.

We get equations (4.3) for $\ell + 1$ with, for $\Omega_{\ell+1}(f_{\ell+1}) = \Omega_{\ell+1}(\omega, z, f_{\ell+1})$ etc.,

$$\begin{aligned} b_{\ell+1}(\omega, z) &= b_{\ell}(\omega, z) + \Omega_{\ell}(\omega, z, \Phi_{\ell+1}), \quad \Omega_{\ell+1}(f_{\ell+1}) = \Omega_{\ell}(f_{\ell+1} + \Phi_{\ell+1}) - \Omega_{\ell}(\Phi_{\ell+1}) \\ c_{\ell+1}(\omega, z) &= c_{\ell}(\omega, z) + \Gamma_{\ell}(\omega, z, \Phi_{\ell+1}), \quad \Gamma_{\ell+1}(f_{\ell+1}) = \Gamma_{\ell}(f_{\ell+1} + \Phi_{\ell+1}) - \Gamma_{\ell}(\Phi_{\ell+1}) \\ d_{\ell+1}(\omega, z) &= d_{\ell}(\omega, z) + Z_{\ell}(\omega, z, \Phi_{\ell+1}), \quad Z_{\ell+1}(f_{\ell+1}) = Z_{\ell}(f_{\ell+1} + \Phi_{\ell+1}) - Z_{\ell}(\Phi_{\ell+1}), \end{aligned}$$

$$(4.14) \quad A_{\ell+1} = P_c(\mathcal{H}_{\omega}) \{ \tilde{A}_{\ell} + c_{\ell+1} \sigma_3 \Phi_{\ell+1} + \mathcal{F}_{\ell}(\omega, z, \Phi_{\ell+1}) - \Phi'_{\ell+1} \cdot (b_{\ell+1}, c_{\ell+1}) \},$$

$$(4.15) \quad \begin{aligned} \mathcal{F}_{\ell+1}(\omega, z, f_{\ell+1}) &= P_c(\mathcal{H}_{\omega}) \{ \mathcal{F}_{\ell}(f_{\ell+1} + \Phi_{\ell+1}) - \mathcal{F}_{\ell}(\Phi_{\ell+1}) \\ &\quad + \Gamma_{\ell+1}(f_{\ell+1}) \sigma_3 \Phi_{\ell} - \Phi'_{\ell+1} \cdot (\Omega_{\ell+1}(f_{\ell+1}), Z_{\ell+1}(f_{\ell+1})) \} + i\omega \partial_{\omega} P_c(\mathcal{H}_{\omega}) f_{\ell+1}. \end{aligned}$$

Notice that $\sigma_1 A_{\ell+1, mn}(\omega) = -A_{\ell+1, nm}(\omega)$ for ω real can be derived from the invariance of (4.3) for $\ell + 1$ solutions of the space where ω is real, z and $\bar{z}$ are complex conjugates and $\sigma_1 f_{\ell+1} = \overline{f_{\ell+1}}$, already stated in Lemma 4.2.

Lemma 4.3. *The following, are analytic functions in $(\omega, z, \bar{z}, f_{\ell+1})$:*

- (1) $b_{\ell+1}, c_{\ell+1}$ and $d_{\ell+1}$ are analytic in $\max\{|\omega - \omega_0|, |z|\} \leq e^{-(2-2^{-\ell-1})}\varepsilon_0$;
- (2) $A_{\ell+1}$ is analytic in $\max\{|\omega - \omega_0|, |z|\} \leq e^{-(2-2^{-\ell-1})}\varepsilon_0$ with values in $\ell^{2,2}$;
- (3) for $\max\{|\omega - \omega_0|, |z|, \|f_{\ell+1}\|_{\ell^{2,-2}}\} \leq e^{-(2-2^{-\ell-1})}\varepsilon_0$, $\Omega_{\ell+1}, \Gamma_{\ell+1}$ and $Z_{\ell+1}$ are analytic;
- (4) we have $\mathcal{F}_{\ell+1}(\omega, z, f_{\ell+1}) = \mathcal{F}(f_{\ell+1}) + \hat{\mathcal{F}}_{\ell+1}(\omega, z, f_{\ell+1})$ with $\mathcal{F}$ the same of induction hypothesis (2) and with $\hat{\mathcal{F}}_{\ell+1}(\omega, z, f_{\ell+1})$ analytic with values in $\ell^{2,2}$ and with domain $\max\{|\omega - \omega_0|, |z|, \|f_{\ell+1}\|_{\ell^{2,-2}}\} \leq e^{-(2-2^{-\ell-1})}\varepsilon_0$.

Proof. (1)–(2) are consequences of $C_A(\ell) D_{\ell} e^{(2^{-\ell-1}-2)\varepsilon_0} < 1$ which we assume, see (4.27). Indeed

$$\|\Phi_{\ell+1}(\omega, z)\|_{\ell^{2,2}} \leq C_A(\ell) D_{\ell} e^{(\ell-1)(2-2^{-\ell})} \varepsilon_0^{-\ell+1} |z|^{\ell+1} \leq C_A(\ell) D_{\ell} e^{(2^{-\ell}-4)\varepsilon_0} \varepsilon_0^2$$

implies $\max\{|\omega - \omega_0|, |z|, \|\Phi_{\ell+1}(\omega, z)\|_{\ell^{2,-2}}\} \leq e^{-(2-2^{-\ell-1})}\varepsilon_0$. For (3), write

$$\|\Phi_{\ell+1}\|_{\ell^{2,-2}} + \|f_{\ell+1}\|_{\ell^{2,-2}} \leq C_A(\ell) D_{\ell} |z|^2 + e^{-(2-2^{-\ell-1})}\varepsilon_0 \leq e^{-(2-2^{-\ell})}\varepsilon_0$$

The last inequality follows from (4.11) and the following inequality, which we assume, see (4.27),

$$(4.16) \quad C_A(\ell) D_{\ell} |z|^2 \leq C_A D_{\ell} \varepsilon_0^2 \leq \varepsilon_0 e^{-2} 2^{-\ell-2}.$$

To prove (4) we proceed similarly by $\mathcal{F}_{\ell+1}(\omega, z, f_{\ell+1}) = \mathcal{F}(f_{\ell+1}) + \widehat{\mathcal{F}}_{\ell+1}(\omega, z, f_{\ell+1})$

$$\begin{aligned}
(4.17) \quad & \widehat{\mathcal{F}}_{\ell+1}(\omega, z, f_{\ell+1}) = P_c(\mathcal{H}_\omega) \widehat{\mathcal{F}}_\ell(\omega, z, f_{\ell+1}) \\
& + \int_{[0,1]^2} P_c(\mathcal{H}_\omega) \mathcal{F}_\ell''(\omega, z, sf_\ell + t\Phi_{\ell+1}) dt ds \cdot f_{\ell+1} \cdot \Phi_{\ell+1} + \\
& P_c(\mathcal{H}_\omega) \{ \Gamma_{\ell+1}(f_{\ell+1}) \sigma_3 \Phi_{\ell+1} - \Phi'_{\ell+1} \cdot (\Omega_{\ell+1}, Z_{\ell+1}) \} + i\dot{\omega} \partial_\omega P_c(\mathcal{H}_\omega) f_{\ell+1}
\end{aligned}$$

Lemma 4.4. *In (4.5) we can choose C_b , C_Ω and C_A in $\ell^\infty(\mathbb{N})$.*

Proof. We have $|\Omega_\ell(\omega, z, \Phi_{\ell+1})| \leq 2C_\Omega(\ell)|z| \|\Phi_{\ell+1}\|_{\ell^{2,2}} \leq 2C_\Omega(\ell)C_A(\ell)D_\ell|z|^3$ for $C_A(\ell)D_\ell\varepsilon_0 < 1$. So for $C_{18}(\ell) := C_b(\ell) + 2\varepsilon_0 C_\Omega(\ell)C_A(\ell)D_\ell$

$$(4.18) \quad |b_{\ell+1}| \leq |b_\ell| + |\Omega_\ell(\omega, z, \Phi_{\ell+1})| \leq C_{18}(\ell)|z|^2.$$

The same bounds hold for $c_{\ell+1}$ and $d_{\ell+1}$. Consider now

$$\begin{aligned}
(4.19) \quad & \Omega_{\ell+1}(\omega, z, f_{\ell+1}) = \Omega_\ell(\omega, z, f_{\ell+1}) \\
& + \int_{[0,1]^2} \Omega_\ell''(\omega, z, sf_{\ell+1} + t\Phi_{\ell+1}(\omega, z)) dt ds \cdot f_{\ell+1} \cdot \Phi_{\ell+1}(\omega, z).
\end{aligned}$$

By induction, by Cauchy integral formula, by (4.11), by (4.16) and by (4.5) we have

$$\begin{aligned}
(4.20) \quad & |\Omega_\ell''(\omega, z, sf_{\ell+1} + t\Phi_{\ell+1}(\omega, z)) \cdot f_{\ell+1} \cdot \Phi_{\ell+1}(\omega, z)| \leq \\
& \leq C_A(\ell)C_\Omega(\ell)D_\ell \frac{|z|^2 \|f_{\ell+1}\|_{\ell^{2,-2}}}{(e^{-2}2^{-\ell-2}\varepsilon_0)^2} (2|z| + \|f_{\ell+1}\|_{\ell^{2,-2}})^2.
\end{aligned}$$

Notice we have used $e^{-2}e^{2^{-\ell}}\varepsilon_0 - e^{-2}e^{2^{-\ell}-2^{-\ell-2}}\varepsilon_0 > \varepsilon_0 e^{-2}2^{-\ell-2}$ and

$$\|sf_{\ell+1} + t\Phi_{\ell+1}\|_{\ell^{2,-2}} \leq \varepsilon_0 e^{-2}2^{-\ell-2} + e^{-(2-2^{-\ell-1})}\varepsilon_0 < e^{-2}e^{2^{-\ell}-2^{-\ell-2}}\varepsilon_0.$$

For $C_{21}(\ell) := C_\Omega(\ell) + e^4 3^4 C_A(\ell)C_\Omega(\ell)2^{2\ell+2}D_\ell\varepsilon_0$

$$\begin{aligned}
(4.21) \quad & |\Omega_{\ell+1}(f_{\ell+1})| \leq |\Omega_\ell(f_{\ell+1})| + e^4 3^4 C_A(\ell)C_\Omega(\ell)2^{2\ell+2}D_\ell\varepsilon_0 |z| \|f_{\ell+1}\|_{\ell^{2,-2}} \\
& \leq C_{21}(\ell) (|z| + \|f_{\ell+1}\|_{\ell^{2,-2}}) \|f_{\ell+1}\|_{\ell^{2,-2}}.
\end{aligned}$$

The same bound holds for $\max\{|\Gamma_{\ell+1}|, |Z_{\ell+1}|, \|\widehat{\mathcal{F}}_\ell\|_{\ell^{2,2}}\}$ by the same argument using

$$(4.22) \quad \|\widehat{\mathcal{F}}_\ell''(\omega, z, sf_\ell + t\Phi_{\ell+1}(\omega, z)) \cdot f_{\ell+1} \cdot \Phi_{\ell+1}(\omega, z)\|_{\ell^{2,2}} \leq \text{rhs}(4.20).$$

Finally we consider $A_{\ell+1}(\omega, z)$. We bound each of the terms in the rhs in (4.14). By Lemma 4.2 for $\Phi_{\ell+1}$ and (4.18) for $c_{\ell+1}$

$$(4.23) \quad \|c_{\ell+1}\sigma_3\Phi_{\ell+1}\|_{\ell^{2,2}} \leq C_{23}(\ell)\varepsilon_0^{-\ell+1}|z|^{\ell+3}, \quad C_{23}(\ell) := C_{18}(\ell)C_A(\ell)D_\ell e^{2(\ell-1)}.$$

Similarly

$$(4.24) \quad \|\widehat{\mathcal{F}}_\ell(\omega, z, \Phi_{\ell+1})\|_{\ell^{2,2}} \leq C_{24}(\ell)\varepsilon_0^{-\ell+1}|z|^{\ell+2}, \quad C_{24}(\ell) := 2C_\Omega(\ell)C_A(\ell)D_\ell e^{2(\ell-1)},$$

$$(4.25) \quad \begin{aligned} \|\Phi'_{\ell+1}(\omega, z) \cdot (b_{\ell+1}, c_{\ell+1})\|_{\ell^{2,2}} &\leq \|\Phi'_{\ell+1}(\omega, z)\|_{\ell^{2,2}}(|b_{\ell+1}| + |c_{\ell+1}|) \\ &\leq 8C_{23}(\ell)\varepsilon_0^{-\ell+1}|z|^{\ell+2}. \end{aligned}$$

We have for $|z| \leq e^{-2+2^{-\ell-1}}\varepsilon_0$

$$(4.26) \quad \begin{aligned} \|\tilde{A}_\ell(\omega, z)\|_{\ell^{2,2}} &\leq \left\| \sum_{m+n \geq \ell+2} \frac{\partial_z^m \partial_{\bar{z}}^n A_\ell(\omega, 0)}{m!n!} z^m \bar{z}^n \right\|_{\ell^{2,2}} \leq \\ &\leq \sum_{j \geq \ell+2} \frac{2^j |z|^j}{j! (\varepsilon_0 \exp(2^{-\ell} - 2))^j} C_A(\ell) (\varepsilon_0 \exp(2^{-\ell} - 2))^2 \\ &\leq C_A(\ell) \sum_{j \geq \ell+2} \frac{2^j |z|^j}{j! (\varepsilon_0 \exp(2^{-\ell} - 2))^{j-2}} \leq C_{26}(\ell) \varepsilon_0^{-\ell} |z|^{\ell+2}, \\ C_{26}(\ell) &:= C_A(\ell) e^2 \frac{2^{\ell+2}}{(\ell+2)! (\exp(2^{-\ell} - 2))^\ell}, \quad C_{26}(1) := \frac{c(1) e^{5/2} 2^3}{3!}. \end{aligned}$$

To close the inequalities we need C_A, C_b, C_Ω in $\ell^\infty(\mathbb{N})$ such that

$$(4.27) \quad \begin{aligned} \varepsilon_0 e^2 \|C_A(j) 2^{j+2} D_j\|_{\ell^\infty} &< 1, \quad \varepsilon_0 \|C_A(j) D_j e^{(2^{-j-1}-2)}\|_{\ell^\infty} < 1, \\ C_b(\ell+1) &\geq C_b(\ell) + \varepsilon_0 C_\Omega(\ell) C_A(\ell) D_\ell, \\ C_\Omega(\ell+1) &\geq C_\Omega(\ell) + \varepsilon_0 C_A(\ell) C_\Omega(\ell) 2^{2\ell+2} e^4 3^2 D_\ell, \\ C_A(\ell+1) &\geq C_A(\ell) \frac{2^{\ell+2} e^{2\ell+2}}{(\ell+2)!} + \\ &+ 8\varepsilon_0^2 [C_b(\ell) + \varepsilon_0 C_\Omega(\ell) C_A(\ell) D_\ell] C_A(\ell) D_\ell e^{2(\ell-1)} + \\ &+ \varepsilon_0 C_\Omega(\ell) C_A(\ell) D_\ell e^{2(\ell-1)}. \end{aligned}$$

We set $C_A(1) = C_b(1) = C_\Omega(1) = c(1)$ and define three sequences recursively using equalities in the last three inequalities in (4.27). They are in $\ell^\infty(\mathbb{N})$ because of the following elementary fact:

Lemma 4.5. *Consider a sequence $x_{n+1} = e^{x_n d_n \varepsilon_0} (x_n + x_n d_n + \varepsilon_0 x_n^N d_n)$ for fixed $N > 1$ and $d \in \ell^1(\mathbb{N})$. Suppose we have inequalities $2\varepsilon_0 x_1 \|d\|_1 \exp(2\|d\|_1) < \log 2$ and $\varepsilon_0 2^{N-1} e^{2(N-1)\|d\|_1} x_1^{N-1} < 1$. Then $x_n \leq 2x_1 \exp(2\|d\|_1)$ for all n .*

Proof. By induction

$$\begin{aligned} x_{n+1} &\leq e^{2\varepsilon_0 x_1 d_n \exp(2\|d\|_1)} \left(1 + d_n + \varepsilon_0 d_n 2^{N-1} e^{2(N-1)\|d\|_1} x_1^{N-1} \right) x_n \\ &\leq e^{2\varepsilon_0 x_1 \|d\|_1 \exp(2\|d\|_1) + 2\|d\|_1} x_1 \leq 2x_1 \exp(2\|d\|_1). \end{aligned}$$

This yields Lemma 4.5. Since we can bound from above our three sequences by a sequence satisfying Lemma 4.5, this also concludes Lemma 4.3.

We define $g(t) = f(t) - \sum_{\ell \geq 2} \Phi_\ell(\omega(t), z(t))$. By $|\omega(t) - \omega_0| + |z(t)| \leq C\epsilon \ll e^{-2}\varepsilon_0$ the series converges. Furthermore $\partial_t g(t) = \partial_t f(t) - \sum_{\ell \geq 2} \partial_t (\Phi_\ell(\omega(t), z(t)))$. We have $g \in \ell_c^2(\mathcal{H}_\omega)$ with $\|g\|_{\ell^2} \lesssim \|f\|_{\ell^2} + |z|^2$ and $\|g\|_{\ell^{2,-2}} \lesssim \|f\|_{\ell^{2,-2}} + |z|^2$. We have particular $f_\ell \rightarrow g$, $\mathcal{H}_\omega f_\ell \rightarrow \mathcal{H}_\omega g$ (notice that $\mathcal{H}_\omega$ is a bounded operator), and $\partial_t f_\ell \rightarrow \partial_t g$ in ℓ^2 . We have $A_\ell(\omega, z) \rightarrow 0$ uniformly for $|\omega - \omega_0| + |z| \leq C\epsilon$. We set $\mathcal{F}_\ell(f_\ell) =$

$$= \mathcal{F}(g) + \widehat{\mathcal{F}}_\ell(g) + (\mathcal{F}(f_\ell) - \mathcal{F}(g)) + \left(\widehat{\mathcal{F}}_\ell(f_\ell) - \widehat{\mathcal{F}}_\ell(g) \right) \rightarrow \mathcal{F}(g) + \lim_{\ell \rightarrow \infty} \widehat{\mathcal{F}}_\ell(g).$$

We have

$$\widehat{\mathcal{F}}_\ell(g) = \widehat{\mathcal{F}}_1(g) + \sum_{j=2}^{\ell} \left(\widehat{\mathcal{F}}_j(g) - \widehat{\mathcal{F}}_{j-1}(g) \right).$$

Since for a fixed C by (4.19) with Ω replaced by $\widehat{\mathcal{F}}$ and by (4.20)

$$\|\widehat{\mathcal{F}}_j(g) - \widehat{\mathcal{F}}_{j-1}(g)\|_{\ell^{2,2}} \leq CD_j 2^{2j+2} e^4 |z|^2 \|g\|_{\ell^{2,-2}} \rightarrow 0 \text{ for } j \nearrow \infty,$$

we have uniform convergence $\widehat{\mathcal{F}}_\ell(g) \rightarrow \widehat{\mathcal{F}}(g)$ in $\ell^{2,2}$. If $|z|_{L_t^\infty} \leq \varepsilon_0$ (recall that $|z|_{L_t^\infty} \leq C\epsilon$) we have $\|\widehat{\mathcal{F}}(\omega, z, g)\|_{\ell^{2,2}} \leq C|z| \|g\|_{\ell^{2,-2}}$. By orbital stability we know $|z| + \|f\|_{\ell^2} \lesssim \epsilon$. Since $\|f - g\|_{\ell^2} \lesssim |z|^2$ we get $\|g\|_{\ell^2} \lesssim \epsilon$. Taking the limit for $\ell \nearrow \infty$ in (4.3) we obtain (4.1). Notice that since $\sigma_1 f_\ell = \overline{f}_\ell$ for all ℓ , we have $\sigma_1 g = \overline{g}$. This ends the proof of Lemma 4.1.

§5 g IS ASYMPTOTICALLY FREE

Our first result is the following one:

Lemma 5.1. *Consider the function g in Lemma 4.1. We have for a fixed C_0*

$$\begin{aligned} \|g\|_{\ell^{\frac{3}{2}r}(\mathbb{Z}, L_t^\infty([n, n+1], \ell^p))} &\leq C_0 \epsilon \text{ for all admissible } (r, p) \\ \|g\|_{L_t^2 \ell^{2, -2}} &\leq C_0 \epsilon. \end{aligned}$$

Proof. Set $P_c(\omega) = P_c(\mathcal{H}_\omega)$. Let ω_0 be as in Theorem 1.2. Set $\vartheta = \omega - \omega_0 + \dot{\gamma}$. We have

$$i\dot{g} - (\mathcal{H}_{\omega_0} + \vartheta P_c(\omega_0) \sigma_3)g = \vartheta [P_c(\omega) - P_c(\omega_0)] \sigma_3 g + \widehat{\mathcal{F}}(\omega, z, g) + O(|g|^7)$$

We split $P_c(\omega_0) = P_+(\omega_0) + P_-(\omega_0)$ with $P_\pm(\omega_0)$ the projections in $\sigma_e(\mathcal{H}_{\omega_0}) \cap \mathbb{R}_\pm$, see §11. Then we rewrite the above equation as

$$\begin{aligned} i\dot{g} - (\mathcal{H}_{\omega_0} + \vartheta(P_+(\omega_0) - P_-(\omega_0))g &= \vartheta [P_c(\omega_0) \sigma_3 - P_+(\omega_0) + P_-(\omega_0)] g \\ &+ \vartheta [P_c(\omega) - P_c(\omega_0)] \sigma_3 g + \widehat{\mathcal{F}}(\omega, z, g) + O(|g|^7). \end{aligned}$$

We prove in Lemma 11.1 that for any pair $s_1, s_2 \in \mathbb{R}$ there is $c_{s_1, s_2}(\omega)$ upper semicontinuous in ω such that for $j = 0, 1$

$$\|P_c(\omega) \sigma_3 - (P_+(\omega) - P_-(\omega))\|_{B(\ell^{2, s_1}, \ell^{2, s_2})} \leq c_{s_1, s_2}(\omega) < \infty.$$

By orbital stability we have $\vartheta = O(\epsilon)$. Then we can write the above equation as

$$i\dot{g} - (\mathcal{H}_{\omega_0} + \vartheta(P_+(\omega_0) - P_-(\omega_0))g = \epsilon O_{loc}(g) + O(|g|^7)$$

for $O_{loc}(g)$ such that $\|O_{loc}(g)\|_{\ell^{2, 2}} \leq C \|g\|_{\ell^{2, -2}}$, for some C . We set

$$(5.1) \quad U_\pm(t, t') = e^{-i(t-t')\mathcal{H}_{\omega_0}} e^{\pm i \int_{t'}^t d\tau (\dot{\gamma}(\tau) + \omega(\tau) - \omega_0)} P_\pm(\omega_0)$$

and we write

$$P_\pm(\omega_0)g(t) = U_\pm(t, 0)g(0) + \int_0^t U_\pm(t, t')(\epsilon O_{loc}(g) + O(|g|^7))dt'.$$

By a standard continuation argument Lemma 5.1 is a consequence of the following result:

Lemma 5.2. *There is a fixed C such that if for all admissible (r, p)*

$$(5.2) \quad \|g\|_{\ell^{\frac{3}{2}r}(\mathbb{Z}, L_t^\infty([n, n+1], \ell^p))} \leq 2C\epsilon, \quad \|g\|_{L_t^2 \ell^{1, -2}} \leq 2C\epsilon$$

then

$$(5.3) \quad \|g\|_{\ell^{\frac{3}{2}r}(\mathbb{Z}, L_t^\infty([n, n+1], \ell^p))} \leq C\epsilon, \quad \|g\|_{L_t^2 \ell^{1, -2}} \leq C\epsilon$$

Proof. Let $\|g\|_{(r,p)} = \|g\|_{\ell^{\frac{3}{2}r}(\mathbb{Z}, L_t^\infty([n, n+1], \ell^p))}$. For fixed $C_{p,\sigma}$

$$\|P_c(\omega) - P_c(\omega_0)\|_{B(\ell^{p,\sigma}, \ell^{p,\sigma})} \leq C_{p,\sigma}|\omega - \omega_0|.$$

So from $C_{p,\sigma}|\omega(t) - \omega_0| \leq 1/2$ for all t and $g(t) = P_c(\omega(t))g(t)$ we conclude $\|g(t)\|_{\ell^{p,\sigma}}/\|P_c(\omega_0)g(t)\|_{\ell^{p,\sigma}} \in [1/2, 2]$. From this we conclude $\|g\|_{(r,p)}/\|P_c(\omega_0)g\|_{(r,p)}$ and $\|g\|_{L_t^2 \ell^{2,-2}}/\|P_c(\omega_0)g\|_{L_t^2 \ell^{2,-2}}$ are in $[1/2, 2]$. Hence it is enough to prove (5.2) $\Rightarrow$ (5.3) with g replaced by $P_c(\omega_0)g$. By Lemmas 3.1-2 we have for fixed constants

$$\|U_\pm(t, 0)g(0)\|_{(r,p)} + \|U_\pm(t, 0)g(0)\|_{L_t^2 \ell^{2,-2}} \lesssim \|g(0)\|_{\ell^2} \lesssim \epsilon.$$

We have by Lemmas 3.5 and 3.6

$$\left\| \int_0^t U_\pm(t, t') \epsilon O_{loc}(g) dt' \right\|_{(r,p) \cap L_t^2 \ell^{2,-2}} \leq C \epsilon \|g\|_{L_t^2 \ell^{2,-2}} = O(\epsilon^2).$$

By Lemma 3.1,

$$\begin{aligned} \left\| \int_0^t U_\pm(t, t') O(|g|^7) dt' \right\|_{(r,p)} &\leq C_0 \|g^7\|_{L_t^1 \ell^2} \leq C_0 \|g\|_{L_t^\infty \ell^2} \|g\|_{L_t^6 \ell^\infty}^6 \\ &\leq C_0 \|g\|_{L_t^\infty \ell^2} \|g\|_{\ell^6(\mathbb{Z}, L_t^\infty([n, n+1]), \ell^\infty)}^6 = O(\epsilon^7). \end{aligned}$$

By (b) Lemma 3.3

$$\left\| \int_0^t U_\pm(t, t') O(|g|^7) dt' \right\|_{L_t^2 \ell^{2,-2}} \leq C \int_0^\infty \|O(g^7)(s)\|_{\ell^2} ds = O(\epsilon^7).$$

This yields Lemma 5.2 and concludes the proof of Lemma 5.1.

Lemma 5.3. *We have ${}^t g(t) = (h(t), \bar{h}(t))$. There exists $r_+ \in \ell^2$ such that $\|r_+\|_{\ell^2} \leq C\epsilon$ for fixed $C = C(\omega_0)$ and*

$$\lim_{t \rightarrow \infty} \|h(t) - e^{it\Delta} r_+\|_{\ell^2} = 0.$$

Proof. First of all, all transformations in §4 preserve the symmetry $\sigma_1 f_\ell = \bar{f}_\ell$. Hence also $\sigma_1 g = \bar{g}$. For $U_\pm(t, t')$ defined in (5.1) we have for $t_1 < t_2$

$$\begin{aligned} &\|U(0, t_2)g(t_2) - U(0, t_1)g(t_1)\|_{\ell^2} \leq \\ &\leq \|U_\pm(0, t')(\epsilon O_{loc}(g) + O(|g|^7))\|_{L^1((t_1, t_2), \ell^2)} \lesssim \\ &\|g\|_{L^2((t_1, t_2), \ell^{2,-2})} + \|g\|_{L^6((t_1, t_2), \ell^\infty)} \rightarrow 0 \text{ for } t_1 \rightarrow \infty. \end{aligned}$$

Then consider $w_+ = P_+(\omega_0)w + P_-(\omega_0)w$ with $P_\pm(\omega_0)w = \lim_{t \rightarrow \infty} U_\pm(0, t)g(t)$. We have

$$\lim_{t \rightarrow \infty} \|P_c(\omega_0)g(t)\|_{\ell^2, -2} = \lim_{t \rightarrow \infty} \|(U_+(t, 0) + U_-(t, 0))P_c(\omega_0)g(0)\|_{\ell^2, -2} = 0$$

with the second equality true for any $g(0) \in \ell^2$. By $\|P_c(\omega_0)g(t)\|_{\ell^2, -2} / \|g(t)\|_{\ell^2, -2} \in [1/2, 2]$ it follows

$$\lim_{t \rightarrow \infty} \|g(t)\|_{\ell^2, -2} = 0$$

Combining the above we have for $\theta(t) = \int_0^t d\tau(\dot{\gamma}(\tau) + \omega(\tau))$

$$\lim_{t \rightarrow \infty} \|g(t) - e^{i[(t\omega_0 - \theta(t) + \theta(0))(P_+(\omega_0) - P_-(\omega_0))]} e^{-it\mathcal{H}_{\omega_0}} w_+\|_{\ell^2} = 0.$$

We claim that the following strong limit exists

$$(5.4) \quad W(\omega_0) = \lim_{t \nearrow \infty} e^{it\mathcal{H}_{\omega_0}} e^{-it(-\Delta + \omega_0)\sigma_3}.$$

The existence of the above limit follows from the existence of the strong limits

$$(5.5) \quad W_1 = \lim_{t \nearrow \infty} e^{it\mathcal{H}_{\omega_0}} e^{-it\mathcal{H}_0} P_c(\mathcal{H}_0) \text{ and } W_2 = \lim_{t \nearrow \infty} e^{it\mathcal{H}_0} e^{it(\Delta - \omega_0)\sigma_3}$$

with $\mathcal{H}_0 = (H + \omega_0)\sigma_3$. The first limit in (5.5) exists by Lemma 9.2. The second limit in (5.5) exists by Pearson's Theorem, see Theorem XI.7[RS], from the fact that $H + \Delta = q$ is trace class. Then the limit in (5.4) exists with $W(\omega_0) = W_1 \circ W_2$ by the "chain rule", Proposition 2 ch.XI [RS]. Furthermore $W(\omega_0)$ is an isomorphism from ℓ^2 to $\ell_c^2(\mathcal{H}_0)$. Set $R_+ = W(\omega_0)^{-1} e^{i\theta(0)(P_+(\omega_0) - P_-(\omega_0))} w_+$. Notice that since $e^{it\omega_0\sigma_3}$ is a unitary matrix periodic in t and $e^{it\omega_0\sigma_3} R_+$ describes a circle in ℓ^2 . Then we have

$$\lim_{t \rightarrow +\infty} \|e^{-it\mathcal{H}_{\omega_0}} W(\omega_0) e^{it\omega_0\sigma_3} R_+ - e^{-it(-\Delta + \omega_0)\sigma_3} e^{it\omega_0\sigma_3} R_+\|_{\ell^2} = 0.$$

Since $W(\omega_0)$ conjugates $\mathcal{H}_{\omega_0}$ into $\sigma_3(-\Delta + \omega_0)$, we get

$$e^{(it\omega_0 + i\theta(0))(P_+(\omega_0) - P_-(\omega_0))} e^{-it\mathcal{H}_{\omega_0}} w_+ = e^{-it\mathcal{H}_{\omega_0}} W(\omega_0) e^{it\omega_0\sigma_3} R_+.$$

Hence the last two limits and the definition of R_+ imply the limit

$$\lim_{t \rightarrow +\infty} \|e^{i\theta(t)\sigma_3} g(t) - e^{it\Delta\sigma_3} R_+\|_{\ell^2} = 0.$$

Hence expressing the last formula in components we obtain Lemma 5.3.

§6 PROOF OF $|z(t)| \approx |z(0)|$

We consider $f = \Phi(\omega, z) + g$. Notice that $\Phi(\omega, z)$ is analytic for $\max\{|\omega - \omega_0|, |z|\} \leq e^{-2}\varepsilon_0$. The equation for z can be written as

$$i\dot{z} - \lambda(\omega)z = a(\omega, z) + Z(\omega, z, g)$$

with $|Z(\omega, z, g)| \leq C|z|\|g\|_{\ell^2, -2}$. We write this equation in the form

$$(6.1) \quad i\dot{z} - \lambda(\omega)z = a(\omega, z) + \langle g, A(\omega, z) \rangle + \langle g, \mathcal{A}(\omega, z, g) \rangle$$

with $\|\mathcal{A}(\omega, z, g)\|_{B(\ell^2, -2, \ell^2, 2)} \leq C$. Our aim is to show:

Lemma 6.1. *There is a fixed constant C such that $|z(0)| \geq C\sqrt{\varepsilon}\|f(0)\|_{\ell^2}$ implies $|z(t)| \approx |z(0)|$ for all t .*

Lemma 6.1 is an immediate consequence of the following lemma:

Lemma 6.2. *There is a change of variables $z = \zeta + \alpha(\omega, \zeta) + \langle g, B(\omega, \zeta) \rangle$ with $|\alpha(\omega, \zeta)| \leq C|\zeta|^2$ and $\langle g, B(\omega, \zeta) \rangle \leq C|\zeta|\|g\|_{\ell^2, -2}$ for a fixed C , such that*

$$(6.2) \quad i\dot{\zeta} - \lambda(\omega)\zeta = d(\omega, |\zeta|^2)\zeta + \langle g, C(\omega, \zeta, g) \rangle$$

with $d(\omega, |\zeta|^2)$ real valued and $|\langle g, C(\omega, \zeta, g) \rangle| \leq C\|g\|_{\ell^2, -2}^2$ for a fixed C .

Assuming Lemma 6.2 we have

$$\frac{d}{dt}|\zeta(t)|^2 = 2\Im [\langle g, C(\omega, \zeta, g) \rangle \bar{\zeta}]$$

and so by Lemma 5.1

$$||\zeta(t)|^2 - |\zeta(0)|^2| \leq C_1\epsilon\|g\|_{\ell^2, -2}^2 \leq C_2\epsilon\|g(0)\|_{\ell^2}^2.$$

Then $|\zeta(0)| \geq \sqrt{2C_2\epsilon}\|g(0)\|_{\ell^2}$ implies $|\zeta(0)|/\sqrt{2} \leq |\zeta(t)| \leq \sqrt{3}|\zeta(0)|$ for all t . Since $|z(t)| \approx |\zeta(t)|$, this concludes Lemma 6.1.

§7 PROOF OF LEMMA 6.2

Lemma 6.1 is a consequence of Lemmas 7.1 and 7.3 below.

Lemma 7.1. *There is a change of variables $z = \varsigma + \alpha(\omega, \varsigma)$ with $|\alpha(\omega, \varsigma)| \leq C|\varsigma|^2$ for a fixed C , such that*

$$(7.1) \quad i\dot{\varsigma} - \lambda(\omega)\varsigma = d(\omega, |\varsigma|^2)\varsigma + \langle g, c(\omega, \varsigma) \rangle + C(\omega, \varsigma, g)$$

with $d(\omega, |\varsigma|^2)$ real, $c(\omega, \varsigma) \in \ell^{2,2}$ and $|C(\omega, \varsigma, g)| \leq C\|g\|_{\ell^{2,-2}}^2$ for a fixed C .

Proof. We write (6.1) in the form

$$(7.2) \quad \begin{aligned} i\dot{z} - \lambda(\omega)z &= \sum_{m+n \geq 2} a_{mn}(\omega)z^m \bar{z}^n + \langle g, A(\omega, z) \rangle + \langle g, \mathcal{A}(\omega, z, g)g \rangle \\ i\dot{\omega} &= b(\omega, z) + \langle g, B(\omega, z) \rangle + \langle g, \mathcal{B}(\omega, z, g)g \rangle. \end{aligned}$$

We define inductively, with (7.3) for $\ell = 1$ equal to (7.2),

$$(7.3) \quad \begin{aligned} i\dot{z}_\ell - \lambda(\omega)z_\ell &= a_\ell(\omega, z_\ell) + \alpha_\ell(\omega, z_\ell, g) \\ i\dot{\omega} &= b_\ell(\omega, z_\ell) + \beta(\omega, z_\ell, g). \end{aligned}$$

Let $\delta_0 = \varepsilon_0^2$, with $\varepsilon_0 > 0$ the constant in §4. We assume the following **inductive hypotheses**.

- (1) $a_\ell, b_\ell, \alpha_\ell$ and β_ℓ are analytic functions in $(\omega, z, \bar{z})$ for $\max\{|\omega - \omega_0|, |z|\} \leq e^{-(2-2^{-\ell})}\delta_0$; if we consider expansions

$$a_\ell(\omega, z) = \sum_{m+n \geq 2} a_{\ell mn}(\omega)z^m \bar{z}^n, \quad b_\ell(\omega, z) = \sum_{m+n \geq 2} b_{\ell mn}(\omega)z^m \bar{z}^n$$

then for ω real, the coefficients $a_{\ell mn}(\omega)$ and $b_{\ell mn}(\omega)$ are real;

- (2) for I_ℓ defined by $(m, n) \in I_\ell$ either if $m + n \geq \ell + 1$ or if $m - n = 1$, we have

$$(7.4) \quad a_\ell(\omega, z) = \sum_{(m,n) \in I_\ell} a_{\ell mn}(\omega)z^m \bar{z}^n, \quad \tilde{a}_\ell(\omega, z) := \sum_{(m,n) \in I_{\ell+1}} a_{\ell mn}(\omega)z^m \bar{z}^n;$$

- (3) the following estimates hold :

$$(7.5) \quad \begin{aligned} |a_\ell(\omega, z_\ell)| &\leq C_a(\ell)|z_\ell|^2, \quad |b_\ell(\omega, z_\ell)| \leq C_b(\ell)|z_\ell|^2 \\ \max\{|\alpha_\ell(\omega, z_\ell, g)|, |\beta_\ell(\omega, z_\ell, g)|\} &\leq C_\alpha(\ell)(|z_\ell| + \|g\|_{\ell^{2,-2}})\|g\|_{\ell^{2,-2}}. \end{aligned}$$

We set $z_{\ell+1} = z_\ell + \phi_{\ell+1}(\omega, z_\ell)$ with

$$\phi_{\ell+1}(\omega, z_\ell) = \sum_{m+n=\ell+1} \frac{a_{\ell mn}(\omega)z_\ell^m \bar{z}_\ell^n}{(m-n-1)\lambda(\omega)} \text{ sum over } m-n \neq 1.$$

Then we get equations (7.3) for $\ell + 1$ with, for $\phi' \cdot (a, b) = \partial_\omega \phi a + \partial_z \phi b - \partial_{\bar{z}} \bar{\phi} \bar{b}$

$$(7.6) \quad \begin{aligned} b_{\ell+1}(\omega, z_{\ell+1}) &= b_\ell(\omega, z_\ell), \quad \beta_{\ell+1}(\omega, z_{\ell+1}, g) = \beta_\ell(\omega, z_\ell, g), \\ a_{\ell+1}(\omega, z_{\ell+1}) &= \tilde{a}_\ell(\omega, z_\ell) - \phi'_{\ell+1}(\omega, z_\ell) \cdot (b_{\ell+1}(\omega, z_{\ell+1}), a_\ell(\omega, z_\ell)), \\ \alpha_{\ell+1}(\omega, z_{\ell+1}, g) &= \alpha_\ell(\omega, z_\ell, g) - \phi'_{\ell+1}(\omega, z_\ell) \cdot (\beta_{\ell+1}(\omega, z_{\ell+1}, g), \alpha_\ell(\omega, z_\ell, g)). \end{aligned}$$

The transformation is designed so that the inductive hypothesis (2) holds for $\ell + 1$, by elementary computation. Similarly, (1) for $\ell + 1$ follows by the definition of $\phi_{\ell+1}(\omega, z_\ell)$, formulas (7.6) and (1) for ℓ . Now we focus on the estimates.

Lemma 7.2. *Estimates (7.5) hold for a fixed C replacing $C_a(\ell)$, $C_b(\ell)$ and $C_\alpha(\ell)$ for all ℓ .*

Proof. Set $\kappa = \sup\{|\lambda^{-1}(\omega)| : |\omega - \omega_0| \leq \varepsilon_0\}$ and $d_\ell := 2^{2\ell+2}\kappa/\ell!$. We have

$$(7.7) \quad \begin{aligned} |\phi_{\ell+1}(\omega, z_\ell)| &\leq \sum_{m+n=\ell+1} \frac{|\lambda^{-1}(\omega)|}{m!n!} |\partial_z^m \partial_{\bar{z}}^n a_\ell(\omega, 0)| |z_\ell|^{\ell+1} \\ &\leq \frac{C_7(\ell) |z_\ell|^{\ell+1}}{(\delta_0 \exp(2^{-\ell} - 2))^{\ell-1}} \leq C_7 |z_\ell|^2, \quad C_7(\ell) := \frac{C_a(\ell) 2^{\ell+1} \kappa}{(\ell+1)!}. \end{aligned}$$

By the Cauchy integral formula for $|\omega - \omega_0| \leq e^{2^{-\ell-1}-2}\delta_0$ and proceeding as for (4.12) and (4.13) we have

$$(7.8) \quad |\phi'_{\ell+1}(\omega, z_\ell)| \leq C_8(\ell) |z_\ell|, \quad C_8(\ell) := C_a(\ell) e^2 d_\ell.$$

By $z_{\ell+1} = z_\ell + \phi_{\ell+1}(\omega, z_\ell)$ we get

$$(7.9) \quad |z_{\ell+1} - z_\ell| \leq C_a(\ell) d_\ell |z_\ell|^2 \leq C_a(\ell) d_\ell \delta_0 |z_\ell|.$$

We assume $C_a(\ell) d_\ell \delta_0 < 1/2 \forall \ell$, see (7.16). Then $|z_{\ell+1} - z_\ell| \leq e^{2C_a(\ell) d_\ell \delta_0} |z_{\ell+1}|$. We have

$$|z_\ell| \leq |z_{\ell+1}| + |z_{\ell+1} - z_\ell| \leq e^{-(2-2^{-\ell-1})} \delta_0 + C_a(\ell) d_\ell e^{-2(2-2^{-\ell-1})} \delta_0^2 < e^{-(2-2^{-\ell})} \delta_0$$

by (4.11) if $C_a(\ell) d_\ell e^{2^{-\ell}-2} 2^{\ell+1} \delta_0 < 1$. We will assume this inequality for the moment, see (7.16). By $(1 - C_a(j) d_j \delta_0)^{-1} = 1 + 2C_a(j) d_j \delta_0$

$$(7.10) \quad |b_{\ell+1}(\omega, z_{\ell+1})| = |b_\ell(\omega, z_\ell)| \leq C_{10}(\ell) |z_{\ell+1}|^2, \quad C_{10}(\ell) := e^{2C_a(\ell) d_\ell \delta_0} C_b(\ell),$$

$$(7.11) \quad |\beta_{\ell+1}(\omega, z_{\ell+1}, g)| = |\beta_\ell(\omega, z_\ell, g)| \leq C_{11}(\ell) (|z_{\ell+1}| + \|g\|_{\ell^2, -2}) \|g\|_{\ell^2, -2}$$

for $C_{11}(\ell) := C_\alpha(\ell) e^{2C_a(\ell) d_\ell \delta_0}$. We have

$$\tilde{a}_\ell(\omega, z_\ell) = a_\ell(\omega, z_\ell) - \hat{a}_\ell(\omega, z_\ell) \text{ with } \hat{a}_\ell(\omega, z) = \sum_{m+n=\ell+1, m-n \neq 1} a_{\ell mn}(\omega) z^m \bar{z}^n.$$

By an analogue of (7.7) we have, by $2C_a(\ell)d_\ell\delta_0 < 1$,

$$(7.12) \quad \begin{aligned} |\widehat{a}_\ell(\omega, z_\ell)| &\leq \frac{C_a(\ell)2^{\ell+1}}{(\ell+1)!(\delta_0 \exp(2^{-\ell} - 2))^{\ell-1}} |z_\ell|^{\ell+1} \leq \\ &\leq \frac{C_a(\ell)2^{\ell+1}}{(\ell+1)!} (1 - C_a(\ell)d_\ell\delta_0)^{-2} |z_{\ell+1}|^2 \leq \frac{C_a(\ell)2^{\ell+3}}{(\ell+1)!} |z_{\ell+1}|^2. \end{aligned}$$

By $a_{\ell+1}(\omega, z_{\ell+1}) = \widetilde{a}_\ell(\omega, z_\ell) - \phi'_{\ell+1}(\omega, z_\ell) \cdot (b_{\ell+1}(\omega, z_{\ell+1}), a_\ell(\omega, z_\ell))$,

$$(7.13) \quad \begin{aligned} |a_{\ell+1}(\omega, z_{\ell+1})| &\leq |a_\ell(\omega, z_\ell)| + \frac{C_a(\ell)2^{\ell+3}}{(\ell+1)!} |z_{\ell+1}|^2 + \\ &+ |\phi'_{\ell+1}(\omega, z_\ell)| |(b_{\ell+1}(\omega, z_{\ell+1}), a_\ell(\omega, z_\ell))| \leq |a_\ell(\omega, z_\ell)| + C_{13}(\ell) |z_{\ell+1}|^2 \\ C_{13}(\ell) &:= \frac{C_a(\ell)2^{\ell+3}}{(\ell+1)!} + \widetilde{C}_{13}(\ell) \text{ for } \ell > 1 \\ \widetilde{C}_{13}(\ell) &:= C_8(\ell)\delta_0 (C_{10}(\ell) + 4C_a(\ell)). \end{aligned}$$

Then

$$(7.14) \quad |a_{\ell+1}(\omega, z_{\ell+1})| \leq C_{14}(\ell) |z_{\ell+1}|^2, \quad C_{14}(\ell) := e^{4C_a(\ell)d_\ell\delta_0} C_a(\ell) + C_{13}(\ell).$$

By $\alpha_{\ell+1}(\omega, z_{\ell+1}, g) = \alpha_\ell(\omega, z_\ell, g) - \phi'_{\ell+1}(\omega, z_\ell) \cdot (\beta_{\ell+1}(\omega, z_{\ell+1}, g), \alpha_\ell(\omega, z_\ell, g))$,

$$(7.15) \quad \begin{aligned} |\alpha_{\ell+1}(\omega, z_{\ell+1}, g)| &\leq C_{15}(\ell) (|z_{\ell+1}| + \|g\|_{\ell^2, -2}) \|g\|_{\ell^2, -2} \\ C_{15}(\ell) &:= e^{4C_a(\ell)d_\ell\delta_0} C_\alpha(\ell) + \delta_0 C_a(\ell) e^2 d_\ell (C_{11}(\ell) + C_\alpha(\ell)). \end{aligned}$$

To close the inequalities we need to find sequences C_a , C_b and C_α in $\ell^\infty(\mathbb{N})$ such that for all ℓ and for $d_\ell := 2^{2\ell+2} \sup_\omega |\lambda^{-1}(\omega)|/\ell!$

$$(7.16) \quad \begin{aligned} C_a(\ell)d_\ell\delta_0 &< 1/2, \quad C_a(\ell)d_\ell e^{2^{-\ell}-2} 2^{\ell+1} \delta_0 < 1 \\ C_a(\ell+1) &\geq C_{14}(\ell), \\ C_b(\ell+1) &\geq C_{10}(\ell), \quad C_\alpha(\ell+1) \geq \max\{C_{11}(\ell), C_{15}(\ell)\}. \end{aligned}$$

For δ_0 small there are such C_a , C_b and C_α in $\ell^\infty(\mathbb{N})$ satisfying (7.16). For example the sequence defined inductively choosing equality in the last two lines in (7.16) and by $C_a(1) = C_b(1) = C_\alpha(1) = c(1)$. This follows from the fact that these sequences are bounded from above by a sequence satisfying Lemma 4.5. This concludes the proof of Lemma 7.2.

The z_ℓ converge to ς . Taking the limit for $\ell \nearrow \infty$ in (7.3) and proceeding as at the end of §4 we see that ς satisfies a system of the form

$$i\dot{\varsigma} - \lambda(\omega)\varsigma = d(\omega, |\varsigma|^2)\varsigma + \gamma(\omega, \varsigma, g)$$

with $|\gamma(\omega, \varsigma, g)| \leq C|\varsigma| \|g\|_{\ell^2, 2}$ and analytic in $(\omega, \varsigma, \bar{\varsigma}, g)$ for $\max\{|\omega - \omega_0|, |\varsigma|\} \leq \delta_0^2$. Expanding $\gamma(\omega, \varsigma, g)$ we get (7.1). This concludes the proof of Lemma 7.1.

Lemma 7.3. *There is a change of variables $\varsigma = \zeta + \langle g, B(\omega, \zeta) \rangle$ with $\langle g, B(\omega, \zeta) \rangle \leq C|\zeta| \|g\|_{\ell^2, -2}$ for a fixed C , such that*

$$(7.17) \quad i\dot{\zeta} - \lambda(\omega)\zeta = d(\omega, |\zeta|^2)\zeta + C(\omega, \varsigma, g)$$

with $d(\omega, |\zeta|^2)$ real valued and $|C(\omega, \varsigma, g)| \leq C\|g\|_{\ell^2, -2}^2$ for a fixed C .

Proof. We write (7.3) in the form

$$(7.18) \quad \begin{aligned} i\dot{\varsigma} - \lambda(\omega)\varsigma &= d(\omega, |\varsigma|^2)\varsigma + \sum_{m+n \geq 1} \langle g, c_{mn}(\omega) \rangle \varsigma^m \bar{\varsigma}^n + \mathcal{C}(\omega, \varsigma, g) \\ i\dot{\omega} &= b(\omega, \varsigma) + \mathcal{B}(\omega, \varsigma, g) \end{aligned}$$

with $|\mathcal{B}(\omega, \varsigma, g)| \leq C\|g\|_{\ell^2, -2}^2$ for a fixed C . We define inductively the following system, which for $\ell = 1$ coincides with (7.18):

$$(7.19) \quad \begin{aligned} i\dot{\varsigma}_\ell - \lambda(\omega)\varsigma_\ell &= d(\omega, |\varsigma_\ell|^2)\varsigma_\ell + \langle g, C_\ell(\omega, \varsigma_\ell) \rangle + \mathcal{C}_\ell(\omega, \varsigma_\ell, g) \\ i\dot{\omega} &= b(\omega, \varsigma_\ell) + \mathcal{B}_\ell(\omega, \varsigma_\ell, g) \\ i\dot{g} - (\mathcal{H}_\omega + \dot{\gamma}P_c(\mathcal{H}_\omega)\sigma_3)g &= G_\ell(\omega, \varsigma_\ell)g + \mathcal{G}_\ell(\omega, \varsigma_\ell, g) + O(|g|^7). \end{aligned}$$

Let $\delta_1 = \delta_0^2$, with δ_0 the constant of Lemmas 7.1–2. We assume the following **inductive hypotheses**.

- (1) $C_\ell(\omega, \varsigma)$ and $B_\ell(\omega, \varsigma)$ are analytic in $(\omega, \varsigma, \bar{\varsigma})$ in $\max\{|\omega - \omega_0|, |\varsigma|\} \leq e^{-(2-2^{-\ell})}\delta_1$ with values in $\ell^{2,2}$; $C_\ell(\omega, \varsigma)$ belongs to $\ell^2(\mathcal{H}_\omega^*)$;
- (2) the following estimates hold:

$$(7.20) \quad \begin{aligned} \|C_\ell(\omega, \varsigma_\ell)\|_{\ell^{2,2}} &\leq C_C(\ell)e^{(\ell-1)(2-2^{-\ell})}\delta_1^{-\ell+1}|\varsigma_\ell|^\ell, \quad |\mathcal{C}_\ell(\omega, \varsigma_\ell, g)| \leq C_C(\ell)\|g\|_{\ell^2, -2}^2, \\ |\mathcal{B}_\ell(\omega, \varsigma_\ell, g)| &\leq C_B(\ell)(|\varsigma_\ell| + \|g\|_{\ell^2, -2})\|g\|_{\ell^2, -2}, \\ \|G_\ell(\omega, \varsigma_\ell)\|_{B(\ell^{2,-2}, \ell^{2,2})} &\leq C_G(\ell)|\varsigma_\ell|, \quad \|\mathcal{G}_\ell(\omega, \varsigma_\ell, g)\|_{\ell^{2,2}} \leq C_G(\ell)\|g\|_{\ell^2, -2}^2. \end{aligned}$$

These hypotheses hold for $\ell = 1$ with $C_C(1) = C_C(1) = C_B(1) = C_G(1) = C_G(1) = c(1)$ for some constant $c(1)$. We expand

$$C_\ell(\omega, \varsigma_\ell) = \sum_{m+n \geq \ell} c_{\ell mn}(\omega) \varsigma_\ell^m \bar{\varsigma}_\ell^n, \quad \tilde{C}_\ell(\omega, \varsigma_\ell) := \sum_{m+n \geq \ell+1} c_{\ell mn}(\omega) \varsigma_\ell^m \bar{\varsigma}_\ell^n.$$

We set $\varsigma_{\ell+1} = \varsigma_\ell + \phi_{\ell+1}(\omega, \varsigma_\ell, g)$

$$\phi_{\ell+1}(\omega, \varsigma_\ell, g) = \langle g, \Phi_{\ell+1}(\omega, \varsigma_\ell) \rangle \text{ with } \Phi_{\ell+1}(\omega, \varsigma_\ell) = \sum_{m+n=\ell} \varsigma_\ell^m \bar{\varsigma}_\ell^n \gamma_{\ell mn}(\omega)$$

$$\gamma_{\ell mn}(\omega) = R_{\mathcal{H}_\omega^*}((n-m+1)\lambda(\omega))c_{\ell mn}(\omega).$$

By induction $C_\ell(\omega, \varsigma_\ell) \in \ell_c^2(\mathcal{H}_\omega^*)$ and so also $c_{\ell mn}(\omega) \in \ell_c^2(\mathcal{H}_\omega^*)$ for all (m, n) . Then $\gamma_{\ell mn}(\omega)$ and $\Phi_{\ell+1}(\omega, \varsigma_\ell) \in \ell_c^2(\mathcal{H}_\omega^*)$. We get equations (7.19) for $\ell + 1$ with:

(7.21)

$$\mathcal{B}_{\ell+1}(\omega, \varsigma_{\ell+1}, g) = \mathcal{B}_\ell(\omega, \varsigma_\ell, g) + b(\omega, \varsigma_\ell) - b(\omega, \varsigma_{\ell+1});$$

(7.22)

$$G_{\ell+1}(\omega, \varsigma_{\ell+1}) = G_\ell(\omega, \varsigma_\ell), \quad \mathcal{G}_{\ell+1}(\omega, \varsigma_{\ell+1}, g) = \mathcal{G}_\ell(\omega, \varsigma_\ell, g);$$

(7.23)

$$\begin{aligned} C_{\ell+1}(\omega, \varsigma_{\ell+1}) &= \tilde{C}_\ell(\omega, \varsigma_{\ell+1}) - [G_\ell(\omega, \varsigma_\ell)]^* \Phi_{\ell+1}(\omega, \varsigma_\ell) - \\ &\quad - \Phi'_{\ell+1}(\omega, \varsigma_\ell) \cdot (b(\omega, \varsigma_\ell), d(\omega, |\varsigma_\ell|^2)_{\varsigma_\ell}) + d^{(1)}(\omega, \varsigma_{\ell+1}); \end{aligned}$$

(7.24)

$$\begin{aligned} \mathcal{C}_{\ell+1}(\omega, \varsigma_{\ell+1}, g) &= \mathcal{C}_\ell(\omega, \varsigma_\ell, g) + \langle g, \tilde{C}_\ell(\omega, \varsigma_\ell) - \tilde{C}_\ell(\omega, \varsigma_{\ell+1}) \rangle \\ &\quad - \langle g, \Phi'_{\ell+1}(\omega, \varsigma_\ell) \cdot (\mathcal{B}_\ell(\omega, \varsigma_\ell, g), \langle g, C_\ell(\omega, \varsigma_\ell) \rangle + \mathcal{C}_\ell(\omega, \varsigma_\ell, g)) \rangle \\ &\quad - \langle \mathcal{G}_\ell(\omega, \varsigma_\ell, g) + O(|g|^7), \Phi_{\ell+1}(\omega, \varsigma_\ell) \rangle + D^{(2)}(\omega, \varsigma_{\ell+1}, g) \end{aligned}$$

with $d^{(1)}(\omega, \varsigma_{\ell+1}) = \partial_g|_{g=0} [d(\omega, |\varsigma_\ell|^2)_{\varsigma_\ell} - d(\omega, |\varsigma_{\ell+1}|^2)_{\varsigma_{\ell+1}}]$ and

$$D^{(2)}(\omega, \varsigma_{\ell+1}, g) = d(\omega, |\varsigma_\ell|^2)_{\varsigma_\ell} - d(\omega, |\varsigma_{\ell+1}|^2)_{\varsigma_{\ell+1}} - \left\langle g, d^{(1)}(\omega, \varsigma_{\ell+1}) \right\rangle.$$

Assumption (1) for $\ell + 1$ holds by an argument analogous to Lemma 4.3. We have

$$\begin{aligned} |\phi_{\ell+1}(\omega, \varsigma_\ell, g)| &\leq \sum_{m+n=\ell} \frac{C_{4.9}}{m!n!} \|\partial_{\varsigma_\ell}^m \partial_{\varsigma_\ell}^n C_\ell(\omega, 0)\|_{\ell^2, 2} |\varsigma_\ell|^\ell \|g\|_{\ell^2, -2} \\ &\leq C_{4.9} |\varsigma_\ell|^\ell \|g\|_{\ell^2, -2} \frac{2^\ell}{\ell!} (e^{2^{-\ell-2}} \delta_1)^{-\ell} \sup_{|z| \leq e^{2^{-\ell-2}} \delta_1} \|C_\ell(\omega, z)\|_{\ell^2, 2} \\ &\leq \frac{C_C(\ell) C_{4.9} 2^\ell}{\ell!} (e^{2^{-\ell-2}} \delta_1)^{-\ell+1} |\varsigma_\ell|^\ell \|g\|_{\ell^2, -2} \text{ for } |\varsigma_\ell| \leq e^{2^{-\ell-2}} \delta_1. \end{aligned}$$

By a similar argument for $|\varsigma_\ell| \leq e^{2^{-\ell-2}} \delta_1$

$$|\partial_{(\varsigma, \bar{\varsigma})} \phi_{\ell+1}(\omega, \varsigma_\ell, g)| \leq \frac{C_C(\ell) C_{4.9} 2^{\ell+2}}{(\ell-1)!} (e^{2^{-\ell-2}} \delta_1)^{-\ell+1} |\varsigma_\ell|^{\ell-1} \|g\|_{\ell^2, -2}$$

and for $|\varsigma_\ell| \leq e^{2^{-\ell-2}} \delta_1$ and $|\omega - \omega_0| \leq e^{2^{-\ell-1}-2} \delta_1$, by an inequality similar to (4.11),

$$|\partial_\omega \phi_{\ell+1}(\omega, \varsigma_\ell, g)| \leq \frac{C_C(\ell) C_{4.9} 2^\ell e^{2^{-\ell-1}}}{\ell! e^{-2} \delta_1} |\varsigma_\ell|^\ell \|g\|_{\ell^2, -2}.$$

Then, for $|\varsigma_\ell| \leq e^{2^{-\ell}-2}\delta_1$, $|\omega - \omega_0| \leq e^{2^{-\ell-1}-2}\delta_1$ and $C_{25}(\ell) := C_C(\ell)C_{4.9}2^{\ell+1}/\ell!$

$$(7.25) \quad \begin{aligned} |\phi_{\ell+1}(\omega, \varsigma_\ell, g)| &\leq C_{25}(\ell)(e^{2^{-\ell}-2}\delta_1)^{-\ell+1}|\varsigma_\ell|^\ell \|g\|_{\ell^2, -2} \leq C_{25}(\ell)|\varsigma_\ell| \|g\|_{\ell^2, -2} \\ |\phi'_{\ell+1}(\omega, \varsigma_\ell, g)| &\leq 2^{12}\ell C_{25}(\ell)(e^{2^{-\ell}-2}\delta_1)^{-\ell+1}|\varsigma_\ell|^{\ell-1} \|g\|_{\ell^2, -2} \leq 2^{12}\ell C_{25}(\ell) \|g\|_{\ell^2, -2}. \end{aligned}$$

We have $\varsigma_{\ell+1} - \varsigma_\ell = \phi_{\ell+1}(\omega, \varsigma_\ell, g)$. So for a fixed C_0 from Lemma 5.1

$$(7.26) \quad \begin{aligned} |\varsigma_{\ell+1} - \varsigma_\ell| &\leq C_{25}(\ell) \|g\|_{\ell^2, -2} |\varsigma_\ell| \leq C_{26}(\ell) \epsilon |\varsigma_\ell| \\ &\leq e^{2C_C(\ell)d_\ell\epsilon} C_C(\ell) d_\ell \epsilon |\varsigma_{\ell+1}|, \quad C_{26}(\ell) := C_0 C_{25}(\ell), \quad d_\ell := C_0 C_{4.9} 2^\ell / \ell!, \end{aligned}$$

where we assume $2C_{26}(\ell)\epsilon = C_C(\ell)d_\ell\epsilon < 1$, see (7.34). We get for $h(\omega, \varsigma) = b(\omega, \varsigma), d(\omega, |\varsigma|)\varsigma$

$$|h(\omega, \varsigma_{\ell+1}) - h(\omega, \varsigma_\ell)| \leq \|D_\varsigma h(\omega, \varsigma)\|_{L^\infty\{|\varsigma| \lesssim |\varsigma_{\ell+1}|\}} |\varsigma_{\ell+1} - \varsigma_\ell|.$$

This yields for $C_{27}(\ell) := e^{2C_C(\ell)d_\ell\epsilon} K(b, d) C_0^{-1} C_C(\ell) d_\ell$ for a fixed $K(b, d)$

$$(7.27) \quad \begin{aligned} |b(\omega, \varsigma_{\ell+1}) - b(\omega, \varsigma_\ell)| &\leq C_{27}(\ell) \delta_1 \|g\|_{\ell^2, -2} |\varsigma_{\ell+1}|, \\ \|d^{(1)}(\omega, \varsigma_{\ell+1})\|_{\ell^2, 2} &\leq C_{27}(\ell) \delta_1 |\varsigma_{\ell+1}|, \\ |D^{(2)}(\omega, \varsigma_{\ell+1}, g)| &\leq C_{27}(\ell) \delta_1 \|g\|_{\ell^2, -2}^2. \end{aligned}$$

$\mathcal{B}_{\ell+1}(\omega, \varsigma_{\ell+1}, g) = \mathcal{B}_\ell(\omega, \varsigma_\ell, g) + b(\omega, \varsigma_\ell) - b(\omega, \varsigma_{\ell+1})$ implies

$$(7.28) \quad \begin{aligned} |\mathcal{B}_{\ell+1}(\omega, \varsigma_{\ell+1}, g)| &\leq C_{28}(\ell) (|\varsigma_{\ell+1}| + \|g\|_{\ell^2, -2}) \|g\|_{\ell^2, -2}, \\ C_{28}(\ell) &= e^{2C_C(\ell)d_\ell\epsilon} C_{\mathcal{B}}(\ell) + C_{27}(\ell) \delta_1. \end{aligned}$$

We bound

$$(7.29) \quad \begin{aligned} \|\tilde{C}_\ell(\omega, \varsigma)\|_{\ell^2, 2} &\leq \sum_{m+n \geq \ell+1} \frac{|\varsigma|^{m+n}}{m!n!} \|\partial_\varsigma^m \partial_{\bar{\varsigma}}^n C_\ell(\omega, 0)\|_{\ell^2, 2} \leq \\ &\leq C_C(\ell) \sum_{j \geq \ell+1} \frac{2^j |\varsigma|^j}{j! (\delta_1 \exp(2^{-\ell} - 2))^{j-1}} \leq \frac{e^2 C_C(\ell) 2^{\ell+1} |\varsigma|^{\ell+1}}{(\ell+1)! (\delta_1 \exp(2^{-\ell} - 2))^\ell}. \end{aligned}$$

By a similar argument for $C_{30}(\ell) = e^2 C_C(\ell) 2^{\ell+2} / \ell!$

$$(7.30) \quad \|\partial_{\varsigma, \bar{\varsigma}} \tilde{C}_\ell(\omega, \varsigma)\|_{\ell^2, 2} \leq C_{30}(\ell) (\delta_1 \exp(2^{-\ell} - 2))^{-\ell} |\varsigma|^\ell.$$

By (7.23) for a constant $K = K(b, d)$ we have

$$\begin{aligned}
(7.31) \quad & \|C_{\ell+1}(\omega, \varsigma_{\ell+1})\|_{\ell^{2,2}} \leq \|\tilde{C}_\ell(\omega, \varsigma_{\ell+1})\|_{\ell^{2,2}} + \|\Phi'_{\ell+1}(\omega, \varsigma_\ell)\|_{\ell^{2,2}} K |\varsigma_\ell|^2 \\
& + \| [G_\ell(\omega, \varsigma_\ell)]^* \|_{B(\ell^{2,2}, \ell^{2,2})} \|\Phi_{\ell+1}(\omega, \varsigma_\ell)\|_{\ell^{2,2}} + \|d^{(1)}(\omega, \varsigma_{\ell+1})\|_{\ell^{2,2}} \\
& \leq C_C(\ell) \left(\frac{e^{2\ell+1}}{(\ell+1)!} |\varsigma_{\ell+1}| + 2^{14} \ell C_{25}(\ell) K \delta_1 |\varsigma_{\ell+1}| \right) \\
& + C_G(\ell) C_{25}(\ell) 2^{\ell+1} \delta_1 |\varsigma_{\ell+1}| + C_{27}(\ell) \delta_1 |\varsigma_{\ell+1}| = C_{31}(\ell) |\varsigma_{\ell+1}|
\end{aligned}$$

with $C_{31}(\ell)$ defined so that the constants match and where we used (7.26) and the assumption $C_0 C_C(\ell) d_\ell \epsilon < 1/2$. By (7.24) we have,

$$\begin{aligned}
& |\mathcal{C}_{\ell+1}(\omega, \varsigma_{\ell+1}, g)| \leq |\mathcal{C}_\ell(\omega, \varsigma_\ell, g)| + \|\tilde{C}_\ell(\omega, \varsigma_\ell) - \tilde{C}_\ell(\omega, \varsigma_{\ell+1})\|_{\ell^{2,2}} \|g\|_{\ell^{2,-2}} + \\
& \|\Phi'_{\ell+1}(\omega, \varsigma_\ell)\|_{\ell^{2,2}} (|\mathcal{B}_\ell(\omega, \varsigma_\ell, g)| + \|C_\ell(\omega, \varsigma_\ell)\|_{\ell^{2,2}} \|g\|_{\ell^{2,-2}} + |\mathcal{C}_\ell(\omega, \varsigma_\ell, g)|) \|g\|_{\ell^{2,-2}} \\
& + \|\Phi_{\ell+1}(\omega, \varsigma_\ell)\|_{\ell^{2,2}} \|\mathcal{G}_\ell(\omega, \varsigma_\ell, g) + O(|g|^7)\|_{\ell^{2,-2}} + |D^{(2)}(\omega, \varsigma_{\ell+1}, g)|.
\end{aligned}$$

We have

$$\begin{aligned}
& \|\tilde{C}_\ell(\omega, \varsigma_\ell) - \tilde{C}_\ell(\omega, \varsigma_{\ell+1})\|_{\ell^{2,2}} \leq C_{31}(\ell) |\varsigma_{\ell+1} - \varsigma_\ell| \leq C_{31}(\ell) C_{25}(\ell) \delta_1 \|g\|_{\ell^{2,-2}}; \\
& \|\Phi'_{\ell+1}(\omega, \varsigma_\ell)\|_{\ell^{2,2}} (|\mathcal{B}_\ell(\omega, \varsigma_\ell, g)| + \|C_\ell(\omega, \varsigma_\ell)\|_{\ell^{2,2}} \|g\|_{\ell^{2,-2}} + |\mathcal{C}_\ell(\omega, \varsigma_\ell, g)|) \\
& \leq 2^{15} \ell C_{25}(\ell) \delta_1 (C_B(\ell) + C_C(\ell) + C_C(\ell)) \|g\|_{\ell^{2,-2}}; \\
& \|\Phi_{\ell+1}(\omega, \varsigma_\ell)\|_{\ell^{2,2}} \|\mathcal{G}_\ell(\omega, \varsigma_\ell, g) + O(|g|^7)\|_{\ell^{2,-2}} \leq C_{25}(\ell) \delta_1 (C_G(\ell) + c_0 \epsilon^5) \|g\|_{\ell^{2,-2}}^2; \\
& |D^{(2)}(\omega, \varsigma_{\ell+1}, g)| \leq C_{27}(\ell) \delta_1 \|g\|_{\ell^{2,-2}}^2.
\end{aligned}$$

So

$$\begin{aligned}
(7.32) \quad & |\mathcal{C}_{\ell+1}(\omega, \varsigma_{\ell+1}, g)| \leq C_{32}(\ell) \|g\|_{\ell^{2,-2}}^2, \quad C_{32}(\ell) := C_C(\ell) + C_{31}(\ell) C_{25}(\ell) \delta_1 + \\
& 2^{15} \ell C_{25}(\ell) \delta_1 (C_B(\ell) + C_C(\ell) + C_C(\ell)) + C_{25}(\ell) \delta_1 (C_G(\ell) + c_0 \epsilon^5) + C_{27}(\ell) \delta_1.
\end{aligned}$$

We have

$$\begin{aligned}
(7.33) \quad & \|G_{\ell+1}(\omega, \varsigma_{\ell+1})\|_{B(\ell^{2,-2}, \ell^{2,2})} = \|G_1(\omega, \varsigma_1)\|_{B(\ell^{2,-2}, \ell^{2,2})} \leq c(1) |\varsigma_1| \\
& \leq C_{33}(\ell) |\varsigma_{\ell+1}|, \quad C_{33}(\ell) := c(1) e^{2\epsilon \|C_C\|_{\infty(j \leq \ell)} \|d_j\|_1} \\
& \|\mathcal{G}_{\ell+1}(\omega, \varsigma_{\ell+1}, g)\|_{\ell^{2,2}} = \|\mathcal{G}_1(\omega, \varsigma_1, g)\|_{\ell^{2,2}} \leq c(1) \|g\|_{\ell^{2,-2}}^2.
\end{aligned}$$

Now we need for all ℓ

$$\begin{aligned}
(7.34) \quad & C_C(\ell) \epsilon d_\ell < 1/2, \quad C_C(\ell) d_\ell e^{2\ell+1} \epsilon < \delta_1 \\
& C_C(\ell+1) \geq C_{31}(\ell), \quad C_B(\ell+1) \geq C_{28}(\ell), \\
& C_C(\ell+1) \geq C_{33}(\ell), \quad C_G(\ell+1) \geq C_{33}(\ell), \quad C_G(\ell+1) = c(1).
\end{aligned}$$

There exist C_C, C_C, C_B and C_G in $\ell^\infty(\mathbb{N})$ satisfying (7.34), see below (7.16). Then we can replace the constants in (7.20) with a fixed constant C . Then $\varsigma_\ell \rightarrow \zeta$ which satisfies the statement of Lemma 7.3.

§8 PROOF OF THEOREM 1.2

We still need to discuss the equation for $\omega(t)$. Recall that we have

$$(8.1) \quad i\dot{\omega} = a(\omega, \zeta) + \langle g, A(\omega, \zeta) \rangle + \mathcal{A}(\omega, \zeta, g)$$

with $|\mathcal{A}(\omega, \zeta, g)| \leq C\|g\|_{\ell^{2,2}}$ and $\|A(\omega, \zeta)\|_{\ell^{2,2}} \leq C|\zeta|$, with $\mathcal{A}(\omega, \zeta, g)$ and $A(\omega, \zeta)$ analytic in $(\omega, \zeta, \bar{\zeta}, g)$. The first step is:

Lemma 8.1. *There is a change of variables $\omega = \varpi + \alpha(\varpi, \zeta) + \langle g, B(\varpi, \zeta) \rangle$ with $|\alpha(\varpi, \zeta)| \leq C|\zeta|^2$ and $\|B(\varpi, \zeta)\|_{\ell^{2,2}} \leq C|\zeta|$ for a fixed C , such that*

$$(8.2) \quad i\dot{\varpi} = D(\varpi, \zeta, g)$$

with $|D(\varpi, \zeta, g)| \leq C\|g\|_{\ell^{2,-2}}^2$ for a fixed C .

An immediate consequence of Lemmas 8.1 and 5.1 is:

Corollary 8.2. *There is a fixed C such that $\|\dot{\varpi}\|_{L^1 \cap L^\infty} < C\epsilon^2$.*

We have:

Corollary 8.3. *For any $\sigma > 0$ there is a fixed C such that for $|z(0)| \geq \epsilon$ and $\|f(0)\|_{\ell^2} \lesssim \epsilon$ we have*

$$\inf_{\kappa, \mu} \|u(t) - e^{i\kappa} \phi_\mu\|_{\ell^{2,-\sigma}} \geq C\epsilon.$$

Proof. We have for ${}^t\xi = (\xi_1, \xi_2)$

$$u(t) = e^{i\theta(t)} \left(\phi_{\omega(t)} + z\xi_1(\omega(t)) + \bar{z}\xi_2(\omega(t)) + A(\omega(t), z(t)) + h(t) \right)$$

with $\|A(\omega(t), z(t))\|_{\ell^{2,2}} \leq C|z(t)|^2$ and $\lim_{t \rightarrow \infty} \|h(t)\|_{\ell^{2,-\sigma}} = 0$. By $\|\xi_2\|_{\ell^{2,2}} \leq C|E_0 - \omega_0|$ and $\|\xi_1 - \varphi_1\|_{\ell^{2,2}} \leq C|E_0 - \omega_0|$, Lemma 2.5, for $t \gg 1$ we have

$$\|e^{-i\theta(t)} u(t) - \phi_{\omega(t)} - z\xi_1(\omega(t))\|_{\ell^{2,-\sigma}} \leq C\epsilon|E_0 - \omega_0|$$

also by $|z| \lesssim \epsilon$, which follows by orbital stability, Lemma 2.3. We have

$$\begin{aligned} \|\phi_\omega + z\xi_1(\omega) - e^{i\kappa} \phi_\mu\|_{\ell^{2,-\sigma}}^2 &\approx \|\phi_\omega + z\xi_1(\omega) - e^{i\kappa} \phi_\mu\|_{\ell^2}^2 \geq \\ &\|\phi_\omega - \phi_\mu\|_{\ell^2}^2 + |z|^2 \|\xi_1(\omega)\|_{\ell^2}^2 - 2|z| |\langle \phi_\omega - \phi_\mu, \xi_1(\omega) \rangle|. \end{aligned}$$

By $\|\xi_1 - \varphi_1\|_{\ell^2} \lesssim |E_0 - \omega_0|$, by $\langle \varphi_0, \varphi_1 \rangle = 0$ and by Lemma 1.1, we have

$$\begin{aligned} \langle \phi_\omega - \phi_\mu, \xi_1(\omega) \rangle &= \left((\omega - E_0)^{\frac{1}{6}} - (\mu - E_0)^{\frac{1}{6}} \right) \langle \varphi_0, O(\omega - E_0) \rangle \\ &+ \langle O((\omega - E_0)^{\frac{7}{6}}) - O((\mu - E_0)^{\frac{7}{6}}), \xi_1(\omega) \rangle = O\left((\omega - \mu)(\omega - E_0)^{\frac{1}{6}}\right). \end{aligned}$$

Hence $\|\phi_\omega + z\xi_1(\omega) - e^{i\kappa}\phi_\mu\|_{\ell^2}^2 \gtrsim |\omega - \mu|^2 + |z|^2 \geq |z|^2$. Then Corollary 8.3 follows from $|z| \gtrsim \epsilon$, Lemma 6.1.

In the rest of the section we prove Lemma 8.1. We have

$$\begin{aligned}
i\dot{\omega} &= a(\omega, \zeta) + \langle g, A(\omega, \zeta) \rangle + \mathcal{A}(\omega, \zeta, g) \\
(8.3) \quad i\dot{\zeta} - \lambda(\omega)\zeta &= d(\omega, |\zeta|^2)\zeta + \mathcal{B}(\omega, \zeta, g) \\
i\dot{g} - (\mathcal{H}_\omega + \dot{\gamma}P_c(\mathcal{H}_\omega)\sigma_3)g &= \widehat{\mathcal{F}}_1(\omega, \zeta)g + O_{loc}(|g|^2) + O(|g|^7).
\end{aligned}$$

We define recursively, where $\omega_1 = \omega$ and (8.4) below for $\ell = 1$ coincides with the equation for ω in (8.3),

$$(8.4) \quad i\dot{\omega}_\ell = a_\ell(\omega, \zeta) + \langle g, A_\ell(\omega, \zeta) \rangle + \mathcal{A}_\ell(\omega, \zeta, g).$$

Let $\delta_2 = \delta_1^2$. We assume:

- (1) for $|\omega - \omega_0| \leq e^{2^{-\ell}-2}\delta_2$ and $|\zeta| \leq \delta_2$ (the equalities below define $\tilde{A}_\ell$ and $\tilde{a}_\ell$)

$$\begin{aligned}
(8.5) \quad a_\ell(\omega, \zeta) &= \sum_{m+n \geq \ell+1} a_{\ell mn}(\omega) \zeta^m \bar{\zeta}^n, \quad \tilde{a}_\ell(\omega, \zeta) := \sum_{m+n \geq \ell+2} a_{\ell mn}(\omega) \zeta^m \bar{\zeta}^n \\
A_\ell(\omega, \zeta) &= \sum_{m+n \geq \ell+1} A_{\ell mn}(\omega) \zeta^m \bar{\zeta}^n, \quad \tilde{A}_\ell(\omega, \zeta) := \sum_{m+n \geq \ell+2} A_{\ell mn}(\omega) \zeta^m \bar{\zeta}^n;
\end{aligned}$$

- (2) the following estimates hold for a fixed C

$$\begin{aligned}
(8.6) \quad |a_\ell(\omega, \zeta)| &\leq C e^{(\ell-1)(2-2^{-\ell})} \delta_2^{-\ell+1} |\zeta|^{\ell+1}, \\
\|A_\ell(\omega, \zeta)\|_{\ell^2, 2} &\leq C e^{(\ell-1)(2-2^{-\ell})} \delta_2^{-\ell+1} |\zeta|^\ell, \quad |\mathcal{A}_\ell(\omega, \zeta, g)| \leq C \|g\|_{\ell^2, -2}^2.
\end{aligned}$$

These facts hold for $\ell = 1$. We set inductively

$$\begin{aligned}
(8.7) \quad \omega_{\ell+1} &= \omega_\ell + \phi_{\ell+1}(\omega, \zeta) + \langle g, \Phi_{\ell+1}(\omega, \zeta) \rangle \\
\phi_{\ell+1}(\omega, \zeta) &= \sum_{m+n=\ell+1, m \neq n} \frac{a_{\ell mn}(\omega) \zeta^m \bar{\zeta}^n}{(m-n)\lambda(\omega)} \\
\Phi_{\ell+1}(\omega, \zeta) &= \sum_{m+n=\ell} \zeta^m \bar{\zeta}^n R_{\mathcal{H}_\omega^*}((n-m)\lambda(\omega)) A_{\ell mn}(\omega).
\end{aligned}$$

$\omega_{\ell+1}$ satisfies (8.4) with

$$\begin{aligned}
(8.8) \quad a_{\ell+1}(\omega, \zeta) &= \tilde{a}_\ell(\omega, \zeta) - \phi'_{\ell+1}(\omega, \zeta)(a(\omega, \zeta), d(\omega, |\zeta|^2)\zeta); \\
\langle g, A_{\ell+1}(\omega, \zeta) \rangle &= \langle g, \tilde{A}_\ell(\omega, \zeta) \rangle - \partial_\omega \phi_{\ell+1}(\omega, \zeta) \langle g, A(\omega, \zeta) \rangle - \\
&\quad - \langle (\dot{\gamma}P_c(\mathcal{H}_\omega)\sigma_3 + \widehat{\mathcal{F}}_1(\omega, \zeta))g, \Phi_{\ell+1}(\omega, \zeta) \rangle; \\
\mathcal{A}_{\ell+1}(\omega, \zeta, g) &= \mathcal{A}_\ell(\omega, \zeta, g) - \phi'_{\ell+1}(\omega, \zeta)(\mathcal{A}(\omega, \zeta, g), \mathcal{B}(\omega, \zeta, g)) - \\
&\quad - \langle O_{loc}(|g|^2) + O(|g|^7), \Phi_{\ell+1}(\omega, \zeta) \rangle.
\end{aligned}$$

More explicitly,

$$(8.9) \quad \begin{aligned} a_{\ell+1}(\omega, \zeta) &= \tilde{a}_\ell(\omega, \zeta) - \partial_\omega \phi_{\ell+1}(\omega, \zeta) a(\omega, \zeta) - \\ &\quad - \partial_\zeta \phi_{\ell+1}(\omega, \zeta) d(\omega, |\zeta|^2) \zeta + \overline{\partial_\zeta \phi_{\ell+1}(\omega, \zeta)} d(\omega, |\zeta|^2) \bar{\zeta} \end{aligned}$$

The estimates can be derived in a manner similar to above proofs and we skip them. We have $\omega_\ell \rightarrow \varpi$ with the latter satisfying an equation of the form (8.2). This completes the proof of Lemma 8.1.

§9 WAVE OPERATORS AND PARTIAL DIAGONALIZATION FOR $\mathcal{H}_\omega$

We set $\mathcal{H}_0 = \sigma_3(H + \omega)$ and write $\mathcal{H}_\omega = \mathcal{H}_0 + B^*(\omega)A$ with $A(\nu) = \langle \nu \rangle^{-\tau}$ with $\tau > 3/2$ and $B^*(\nu, \omega)$ a C^2 function in ω with values in the space of 2×2 real valued matrices. For any ω in some compact set K there is a constant $c(K) > 0$ and $\alpha > 0$ such that $|e^{\alpha|\nu|} B^*(\nu, \omega)| \leq c_m(K)(\omega - E_0) \forall \nu \in \mathbb{Z}$.

Lemma 9.1. *For $\tau > 1$ there exists $C = C(\tau, \omega)$ such that for all $z \in \mathbb{C} \setminus \sigma_e(\mathcal{H}_\omega)$*

$$(9.1) \quad \|R_{\mathcal{H}_\omega}(z)P_c(\mathcal{H}_\omega)\|_{B(\ell^{2,\tau}, \ell^{2,-\tau})} \leq C.$$

There is a neighborhood U of $\sigma_e(\mathcal{H}_\omega)$ in $\mathbb{C}$ such that for any $z \in U \setminus \sigma_e(\mathcal{H}_\omega)$

$$(9.2) \quad \|R_{\mathcal{H}_\omega}(z)\|_{B(\ell^{2,\tau}, \ell^{2,-\tau})} \leq C.$$

The following limits are well defined

$$(9.3) \quad \lim_{\epsilon \rightarrow 0^+} R_{\mathcal{H}_\omega}(\lambda \pm i\epsilon) = R_{\mathcal{H}_\omega}^\pm(\lambda) \text{ in } C^0(\sigma_e(\mathcal{H}_\omega), B(\ell^{2,\tau}, \ell^{2,-\tau})).$$

Proof. First of all (9.2) implies (9.1). By Lemma 5.7 [CT] we have

$$(1) \quad \|\langle x \rangle^{-\tau} R_H(z, \cdot, \cdot) \langle y \rangle^{-\tau}\|_{\ell^2(\mathbb{Z}^2)} \leq C \text{ for } z \text{ close to } [0, 4]$$

with the following limits well defined

$$(2) \quad \lim_{\epsilon \rightarrow 0^+} R_H(\lambda \pm i\epsilon) = R_H^\pm(\lambda) \text{ in } C^0([0, 4], B(\ell^{2,\tau}, \ell^{2,-\tau})).$$

This implies

$$(3) \quad \|\langle x \rangle^{-\tau} R_{\mathcal{H}_0}(z, \cdot, \cdot) \langle y \rangle^{-\tau}\|_{\ell^2(\mathbb{Z}^2)} \leq C \text{ for } z \text{ close to } \sigma_e(\mathcal{H}_\omega)$$

with the following limits well defined

$$(4) \quad \lim_{\epsilon \rightarrow 0^+} R_{\mathcal{H}_0}(\lambda \pm i\epsilon) = R_{\mathcal{H}_0}^\pm(\lambda) \text{ in } C^0(\sigma_e(\mathcal{H}_\omega), B(\ell^{2,\tau}, \ell^{2,-\tau})).$$

For $A = \langle x \rangle^{-\tau}$ we write

$$(5) \quad AR_{\mathcal{H}_\omega}(z) = (1 + AR_{\mathcal{H}_0}(z)B^*)^{-1}AR_{\mathcal{H}_0}(z).$$

(3) implies $\|AR_{\mathcal{H}_0}(z)B^*\|_{B(\ell^2, \ell^2)} \leq C(\omega - E_0) \ll 1$. (3)–(4) imply (9.2)–(9.3). This concludes the proof of Lemma 9.1.

We have:

Lemma 9.2. *We have:*

- (A) $\mathcal{H}_\omega$ does not have resonances at $\pm\omega$ and at $\pm(4+\omega)$.
- (B) $\sigma_e(\mathcal{H}_\omega)$ does not contain eigenvalues.
- (C) There are isomorphisms inverses of each other $W(\omega) \in B(\ell_c^2(\mathcal{H}_0), \ell_c^2(\mathcal{H}_\omega))$ and $Z(\omega) \in B(\ell_c^2(\mathcal{H}_\omega), \ell_c^2(\mathcal{H}_0))$, defined as follows: for $u \in \ell_c^2(\mathcal{H}_0)$, and v such that $\sigma_3 v \in \ell_c^2(\mathcal{H}_\omega)$,

$$\langle Wu, v \rangle = \langle u, v \rangle + \lim_{\varepsilon \rightarrow 0^+} \frac{1}{2\pi i} \int_{-\infty}^{+\infty} \langle AR_{\mathcal{H}_0}(\lambda + i\varepsilon)u, BR_{\mathcal{H}_\omega^*}(\lambda + i\varepsilon)v \rangle d\lambda;$$

for $u \in \ell_c^2(\mathcal{H}_\omega)$, $v \in \ell_c^2(\mathcal{H}_0)$,

$$\langle Zu, v \rangle = \langle u, v \rangle + \lim_{\varepsilon \rightarrow 0^+} \frac{1}{2\pi i} \int_{-\infty}^{+\infty} \langle AR_{\mathcal{H}_\omega}(\lambda + i\varepsilon)u, BR_{\mathcal{H}_0}(\lambda + i\varepsilon)v \rangle d\lambda.$$

Then $P_c(H_\omega)\mathcal{H}_\omega = W\mathcal{H}_0Z$. $\|W(\omega)\|_{B(\ell_c^2(\mathcal{H}_0), \ell_c^2(\mathcal{H}_\omega))}$ and $\|Z(\omega)\|_{B(\ell_c^2(\mathcal{H}_\omega), \ell_c^2(\mathcal{H}_0))}$ are uniformly locally bounded in ω .

Proof. (A) and (B) follow by standard arguments from the fact that V_ω is small. (C) follows from (1)–(4) below. Specifically we need to show that there is a fixed $c > 0$ such that $\forall \varepsilon \neq 0$

- (1) $\int \|\langle x \rangle^{-\tau} R_{\mathcal{H}_0}(\lambda + i\varepsilon)u\|_{\ell^2}^2 d\lambda \leq c\|u\|_{\ell^2}^2$ for all $u \in \ell_c^2(\mathcal{H}_0)$
- (2) $\int \|BR_{\mathcal{H}_0}(\lambda + i\varepsilon)u\|_{\ell^2}^2 d\lambda \leq c\|u\|_2^2$ for all $u \in \ell_c^2(\mathcal{H}_0)$
- (3) $\int \|BR_{\mathcal{H}_\omega^*}(\lambda + i\varepsilon)u\|_{\ell^2}^2 d\lambda \leq c\|u\|_{\ell^2}^2$ for all $u \in \ell^2(\mathcal{H}_\omega^*) := \sigma_3 \ell^2(\mathcal{H}_\omega)$
- (4) $\int \|\langle x \rangle^{-\tau} R_{\mathcal{H}_\omega}(\lambda + i\varepsilon)u\|_{\ell^2}^2 d\lambda \leq c\|u\|_{\ell^2}^2$ for all $u \in \ell_c^2(\mathcal{H}_\omega)$.

(1)–(4) are consequences of (9.1) and of inequalities (2)–(3) in Lemma 9.1.

We have:

Lemma 9.3. *For any $u \in \ell^2$ we have*

$$(1) \quad P_c(\mathcal{H}_\omega)u = \lim_{\varepsilon \rightarrow 0^+} \frac{1}{2\pi i} \int_{\sigma_e(\mathcal{H}_\omega)} [R_{\mathcal{H}_\omega}(\lambda + i\varepsilon) - R_{\mathcal{H}_\omega}(\lambda - i\varepsilon)] u d\lambda.$$

Proof. By $\sigma_e(\mathcal{H}_\omega) = \sigma_e(\mathcal{H}_0) = [\omega, 4+\omega] \cup [-4-\omega, -\omega]$ and by the spectral theorem we have

$$(2) \quad P_c(\mathcal{H}_0)v = \lim_{\varepsilon \rightarrow 0^+} \frac{1}{2\pi i} \int_{\sigma_e(\mathcal{H}_\omega)} [R_{\mathcal{H}_0}(\lambda + i\varepsilon) - R_{\mathcal{H}_0}(\lambda - i\varepsilon)] v d\lambda.$$

To prove (1) in Lemma 9.3 we observe that for $u \in \ell_d^2(\mathcal{H}_\omega)$ both sides of (1) are 0. Hence it is enough to prove (1) for $u \in \ell_c^2(\mathcal{H}_\omega)$. Then $u = W(\omega)v$ with $v \in \ell_c^2(\mathcal{H}_0)$. Then (1) $u \in \ell_c^2(\mathcal{H}_\omega)$ can be rewritten as

$$\begin{aligned} u &= W(\omega)v = W(\omega) \lim_{\epsilon \rightarrow 0^+} \frac{1}{2\pi i} \int_{\sigma_e(\mathcal{H}_\omega)} [R_{\mathcal{H}_0}(\lambda + i\epsilon) - R_{\mathcal{H}_0}(\lambda - i\epsilon)] v d\lambda \\ &= \lim_{\epsilon \rightarrow 0^+} \frac{1}{2\pi i} \int_{\sigma_e(\mathcal{H}_\omega)} [R_{\mathcal{H}_\omega}(\lambda + i\epsilon) - R_{\mathcal{H}_\omega}(\lambda - i\epsilon)] W(\omega)v d\lambda. \end{aligned}$$

This concludes Lemma 9.3.

Finally, we obtain the limiting absorption principle:

Lemma 9.4. *For any $u \in \ell^{2,\tau}$ with $\tau > 3/2$ we have*

$$P_c(\mathcal{H}_\omega)u = \frac{1}{2\pi i} \int_{\sigma_e(\mathcal{H}_\omega)} [R_{\mathcal{H}_\omega}^+(\lambda) - R_{\mathcal{H}_\omega}^-(\lambda)] u d\lambda.$$

Proof. For $u \in \ell^{2,\tau}$ the $\epsilon \rightarrow 0^+$ limit in (1) Lemma 9.4 converges in $\ell^{2,-\tau}$ to the integral in Lemma 9.5.

§10 DISPERSIVE THEORY FOR $\mathcal{H}_\omega$: PROOF OF LEMMA 3.1

Lemma 9.2 implies $\|P_c(\mathcal{H}_\omega)e^{it\mathcal{H}_\omega}\|_{B(\ell^2,\ell^2)} \leq C$ for a fixed $C > 0$ and yields Lemma 3.1 for $p = 2$. By interpolation the rest of Lemma 3.1 will be a consequence of case $p = 1$. Lemma 9.4 implies for $u \in \mathcal{S}(\mathbb{Z})$

$$P_c(\mathcal{H}_\omega)e^{it\mathcal{H}_\omega}u = \frac{1}{2\pi i} \int_{\sigma_e(\mathcal{H}_\omega)} e^{i\lambda t} [R_{\mathcal{H}_\omega}^+(\lambda) - R_{\mathcal{H}_\omega}^-(\lambda)] u d\lambda.$$

We expand $R_{\mathcal{H}_\omega}^\pm(\lambda) = \sum_{j=0}^\infty (-1)^j (R_{\mathcal{H}_0}^\pm(\lambda)V_\omega)^j R_{\mathcal{H}_0}^\pm(\lambda)$. Correspondingly

$$\begin{aligned} P_c(\mathcal{H}_\omega)e^{it\mathcal{H}_\omega}u &= \sum_{j=1}^\infty (-1)^j \frac{1}{2\pi i} \int_{\sigma_e(\mathcal{H}_\omega)} e^{i\lambda t} (R_{\mathcal{H}_0}^+(\lambda)V_\omega)^j R_{\mathcal{H}_0}^+(\lambda) u d\lambda \\ &\quad - \sum_{j=1}^\infty (-1)^j \frac{1}{2\pi i} \int_{\sigma_e(\mathcal{H}_\omega)} e^{i\lambda t} (R_{\mathcal{H}_0}^-(\lambda)V_\omega)^j R_{\mathcal{H}_0}^-(\lambda) u d\lambda + P_c(\mathcal{H}_0)e^{it\mathcal{H}_0}u. \end{aligned}$$

We have $\|P_c(\mathcal{H}_0)e^{it\mathcal{H}_0}\|_{B(\ell^1,\ell^\infty)} \leq C\langle t \rangle^{-\frac{1}{3}}$ by [CT,PS]. We consider now a generic term in the above summations. By $\sigma_e(\mathcal{H}_\omega) = [\omega, 4 + \omega] \cup [-4 - \omega, -\omega]$ it is not restrictive to focus on what follows, for $K(E) = (R_{\mathcal{H}_0}^+(E + \omega)V_\omega)^j$ with $j \geq 0$,

$$Tu = \int_0^4 e^{iEt} R_{\mathcal{H}_0}^+(E + \omega)V_\omega K(E) R_{\mathcal{H}_0}^+(E + \omega) u dE.$$

Let $P_+^0 = \text{diag}(1, 0)$ and $P_-^0 = \text{diag}(0, 1)$. Then $T = P_+^0 T P_+^0 + P_-^0 T P_+^0 + P_+^0 T P_-^0 + P_-^0 T P_-^0$. We consider separately these four operators. We focus on $T_{++} = P_+^0 T P_+^0$. We have

$$T_{++}(\mu, \nu) = \sum_{(\nu', \mu') \in \mathbb{Z}^2} \int_0^4 e^{i\lambda t} P_+^0 R_H^+(\lambda, \nu, \nu') V_\omega(\nu') K(\lambda, \nu', \mu') R_H^+(\lambda, \mu', \mu) P_+^0 d\lambda.$$

Recall now, see for instance Lemma 5.9 [CT], that

$$(10.1) \quad \begin{aligned} R_H^+(\lambda, \nu, \nu') &= -\frac{f_+(\nu, \theta) f_-(\nu', \theta)}{W(\theta)} \quad \text{for } \nu \geq \nu', \\ R_H^+(\lambda, \nu, \nu') &= -\frac{f_-(\nu, \theta) f_+(\nu', \theta)}{W(\theta)} \quad \text{for } \nu < \nu' \end{aligned}$$

where $\theta \in [0, \pi]$ is such that $\lambda = 2(1 - \cos \theta)$, where $f_\pm(\nu, \theta)$ are the Jost functions satisfying

$$H f_\pm(\nu, \theta) = \lambda f_\pm(\nu, \theta) \text{ with } \lim_{\nu \rightarrow \pm\infty} [f_\pm(\nu, \theta) - e^{\mp i\nu\theta}] = 0$$

and where $W(\theta) = f_+(\nu+1, \theta) f_-(\nu, \theta) - f_+(\nu, \theta) f_-(\nu+1, \theta)$ is the Wronskian. We recall that $W(\theta)$ is C^∞ in $\mathbb{R}/2\pi\mathbb{Z}$ and that $W(\theta) \neq 0$ for all θ , see Lemmas 5.3 and 5.5 in [CT]. We split the above sum in various terms

$$(10.2) \quad T_{++}(\mu, \nu) = \sum_{\nu' \leq \nu, \mu \leq \mu'} \cdots + \sum_{\nu' > \nu, \mu \leq \mu'} \cdots + \sum_{\nu' \leq \nu, \mu > \mu'} \cdots + \sum_{\nu' > \nu, \mu > \mu'} \cdots.$$

Let $m_\pm(\nu, \theta) = e^{\pm i\nu\theta} f_\pm(\nu, \theta)$. Then the first term in the rhs in (10.1) is

$$(10.3) \quad \sum_{\nu' \leq \nu, \mu \leq \mu'} \int_0^\pi e^{it(2-2\cos\theta) + i(\nu-\mu)\theta} m_-(\nu, \theta) A(\theta, \nu', \mu') m_+(\mu, \theta) \sin \theta d\theta$$

with $A(\theta, \nu', \mu') = f_+(\nu', \theta) V_\omega(\nu') K(\lambda, \nu', \mu') f_-(\mu', \theta)$. We have

$$\sup_{\nu' \leq \nu, \mu \leq \mu'} (\langle \nu' \rangle^2 \langle \mu' \rangle^2 \|m_-(\nu, \theta) A(\theta, \nu', \mu') m_+(\mu, \theta)\|_{W^{1,1}(0, \pi)}) \leq C^j (\omega - E_0)^{j+1}.$$

Then by stationary phase $|(10.3)| \leq C^j (\omega - E_0)^{j+1} \langle t \rangle^{-\frac{1}{3}}$. The other terms in the rhs of (10.2) can be bounded using

$$f_\mp(\nu, \theta) = \frac{1}{T(\theta)} \overline{f_\pm(\nu, \theta)} + \frac{R_\pm(\theta)}{T(\theta)} f_\pm(\nu, \theta)$$

where transmission and reflection coefficients are defined using the Wronskians

$$T(\theta) = \frac{\pm 2i \sin \theta}{[f_\mp(\theta), f_\pm(\theta)]}, \quad R_\pm(\theta) = -\frac{[f_\mp(\theta), \overline{f_\pm(\theta)}]}{[f_\mp(\theta), f_\pm(\theta)]},$$

see [CT]. Finally, the remaining terms $P_-^0 T P_+^0$, $P_+^0 T P_-^0$ and $P_-^0 T P_-^0$ can be bounded similarly. So we have $\|T\|_{B(\ell^1, \ell^\infty)} \leq C^j (\omega - E_0)^{j+1} \langle t \rangle^{-\frac{1}{3}}$. By summing up on j and for $C(\omega - E_0) < 1$ we obtain $\|P_c(\mathcal{H}_\omega) e^{it\mathcal{H}_\omega}\|_{B(\ell^1, \ell^\infty)} \leq C \langle t \rangle^{-\frac{1}{3}}$.

§11 THE PROJECTIONS $P_{\pm}(\omega)$

Let $u \in \mathcal{S}(\mathbb{Z})$. We set

$$\begin{aligned} P_{\pm}(\omega)u &= \frac{1}{2\pi i} \int_{\sigma_e(\mathcal{H}_{\omega}) \cap \mathbb{R}_{\pm}} [R_{\mathcal{H}_{\omega}}^+(\lambda) - R_{\mathcal{H}_{\omega}}^-(\lambda)] u d\lambda \\ &= \lim_{\epsilon \rightarrow 0^+} \frac{1}{2\pi i} \int_{\sigma_e(\mathcal{H}_{\omega}) \cap \mathbb{R}_{\pm}} [R_{\mathcal{H}_{\omega}}(\lambda + i\epsilon) - R_{\mathcal{H}_{\omega}}(\lambda - i\epsilon)] u d\lambda. \end{aligned}$$

Notice that we have $P_{\pm}(\omega) = Z(\omega)P_{\pm}^0 P_c(H)W(\omega)$. Hence $P_{\pm}(\omega)$ extend into projections in ℓ^2 . During the course of the proof of Lemma 5.1 we used the following fact, which we prove:

Lemma 11.1. *For any pair $s_1, s_2 \in \mathbb{R}$ there is $c_{s_1, s_2}(\omega)$ upper semicontinuous in ω such that for $j = 0, 1$*

$$(1) \quad \|P_c(\omega)\sigma_3 - (P_+(\omega) - P_-(\omega))\|_{B(\ell^{2, s_1}, \ell^{2, s_2})} \leq c_{s_1, s_2}(\omega) < \infty.$$

Proof. For this proof we set $\mathcal{H} = \mathcal{H}_{\omega}$, $\mathcal{H}_0 = \sigma_3(H + \omega)$, $R_0(z) = (\mathcal{H}_0 - z)^{-1}$ and $R(z) = (\mathcal{H} - z)^{-1}$. To prove (1) it is enough to write $P_c = P_+ + P_-$ and to prove $\|[P_{\pm}\sigma_3 \mp P_{\pm}]g\|_{\ell^{2, M}} \leq c\|g\|_{\ell^{2, -N}}$. It is not restrictive to consider only P_+ . Setting $H = H_0 + V$, we write

$$(2) \quad \sum_{\pm} \pm R(\lambda \pm i\epsilon) = \sum_{\pm} \pm (1 + R_0(\lambda \pm i\epsilon)V)^{-1} R_0(\lambda \pm i\epsilon).$$

By elementary computation

$$R_0(\lambda \pm i\epsilon)\sigma_3 = R_0(\lambda \pm i\epsilon) - 2(H + \omega + \lambda \pm i\epsilon)^{-1} \text{diag}(0, 1).$$

Therefore

$$\text{rhs}(2)\sigma_3 = \text{rhs}(4) + 2 \sum_{\pm} \pm (1 + R_0(\lambda \pm i\epsilon)V)^{-1} \text{diag}(0, 1) R_H(-\omega - \lambda \mp i\epsilon).$$

Hence we are reduced to show that $Ku =$

$$\lim_{\epsilon \rightarrow 0^+} \sum_{\pm} \pm \int_{\mathbb{R}_+ \cap \sigma_c(\mathcal{H}_0)} (1 + R_0(\lambda \pm i\epsilon)V)^{-1} \text{diag}(0, 1) R_H(-\omega - \lambda \mp i\epsilon) u d\lambda$$

defines an operator such that for some fixed c

$$(3) \quad \|Ku\|_{\ell^{2, M}} \leq c\|u\|_{\ell^{2, -N}}$$

We expand $(1 + R_0 V)^{-1} = \sum_{j=0}^{\infty} [-R_0 V]^j$ and we consider the corresponding decomposition $K = \sum_{j=0}^{\infty} K_j^0$. We have $K_0^0 = 0$ since for any $u \in \ell^2$ we have

$$\lim_{\epsilon \rightarrow 0^+} \int_{\mathbb{R}_+ \cap \sigma_c(\mathcal{H}_0)} \sum_{\pm} \pm (H + \omega + \lambda \pm i\epsilon)^{-1} \text{diag}(0, 1) u d\lambda = 0.$$

We next consider K_j^0 for $j > 0$ and prove

$$(4) \quad \|K_j^0 u\|_{\ell^2, M} \leq c^j (\omega - E_0)^j \|u\|_{\ell^2, -N} \text{ for a fixed } c.$$

We have that for γ a closed path around $[\omega, 4 + \omega]$ we have

$$K_j^0 u = \int_{\gamma} R_0(z) V(R_0(z) V)^{j-1} \text{diag}(0, 1) R_H(-\omega - z) u dz.$$

We have $\|V(R_0(z) V)^{j-1}\|_{B(\ell^2, -N, \ell^2, M)} \leq \tilde{c}^j (\omega - E_0)^j$. For $z \in \gamma$ we have

$$|R_0(z, \nu, \mu)| + |R_H(-\omega - z, \nu, \mu)| \leq \beta e^{-\alpha|\nu - \mu|}$$

for some fixed $\alpha > 0$ and $\beta > 0$. Then

$$\begin{aligned} & \|K_j^0\|_{B(\ell^2, -N, \ell^2, M)} \\ & \leq \beta^2 \|e^{-\alpha|\cdot|} * \|_{B(\ell^2, -N, \ell^2, -N)} \|V(R_0(z) V)^{j-1}\|_{B(\ell^2, -N, \ell^2, M)} \|e^{-\alpha|\cdot|} * \|_{B(\ell^2, -N, \ell^2, M)} \\ & \leq c^j (\omega - E_0)^j. \end{aligned}$$

This yields Lemma 11.1.

§APPENDIX A: EXISTENCE OF H SATISFYING (H1)–(H3)

Lemmas A.1 and A.2 below, together prove the existence of H satisfying (H1)–(H3). Lemma A.1 uses standard perturbation arguments. Lemma A.2 proves that, in some sense, 0 average potentials generically satisfy the second condition in (1) below. We recall that the resolvent $R_{-\Delta}(z)$ for $z \in \mathbb{C} \setminus [0, 4]$ has kernel

$$R_{-\Delta}(\mu, \nu, z) = \frac{1}{2i \sin \theta} e^{-i\theta|\mu - \nu|}, \quad \mu, \nu \in \mathbb{Z},$$

with θ a solution to $2(1 - \cos \theta) = z$ with $\Im \theta \leq 0$. We will consider $q(\mu) \in S(\mathbb{Z})$.

Lemma A.1. *Suppose that for some $a_0 > 0$ and C_0 we have $|q(\mu)| \leq C_0 e^{-a_0|\mu|}$ and that the following two conditions are satisfied:*

$$(1) \quad \sum_{\mu \in \mathbb{Z}} q(\mu) = 0, \quad \sum_{\mu, \nu \in \mathbb{Z}} |\mu - \nu| q(\mu) q(\nu) < 0.$$

Then there is $\varepsilon_0 > 0$ such that for any $0 < \varepsilon < \varepsilon_0$ the operator $H = -\Delta + \varepsilon q$ satisfies hypotheses (H1)–(H3).

Proof. For $z \notin \sigma(H)$ we have $R_H(z) = (1 + R_{-\Delta}(z)\varepsilon q)^{-1} R_{-\Delta}(z)$. So there are two eigenvalues, one in $(-\infty, 0)$ and the other in $(4, \infty)$, exactly if $(1 + R_{-\Delta}(z)\varepsilon q)^{-1}$ is singular in two such points. By Fredholm theory the singular points occur in correspondence to values of z such that $\ker(1 + R_{-\Delta}(z)\varepsilon q) \neq 0$. If we set $b(\nu) = \sqrt{|q(\nu)|}$ and $a(\nu) = b(\nu)\text{sign } q(\nu)$ we have $q(\nu) = a(\nu)b(\nu)$. By standard arguments the map $u \rightarrow v = bu$ establishes an isomorphism $\ker(1 + R_{-\Delta}(z)\varepsilon q) \rightarrow \ker(1 + bR_{-\Delta}(z)\varepsilon a)$ with inverse $u = -R_{-\Delta}(z)av$. So $\ker(1 + R_{-\Delta}(z)\varepsilon q) \neq 0$ exactly if $\ker(1 + bR_{-\Delta}(z)\varepsilon a) \neq 0$. We split the analysis in two parts. We first look at z near 0. We will later look at z near 4. For z near 0 write

$$R_{-\Delta}(\mu, \nu, z) = \frac{1}{2i \sin \theta} - D_0(\mu, \nu, z), \quad D_0(\mu, \nu, z) = \frac{1 - e^{-i\theta|\mu-\nu|}}{2i \sin \theta}.$$

Now we have

$$\begin{aligned} 1 + bR_{-\Delta}(z)\varepsilon a &= 1 + \varepsilon \frac{b\langle \cdot, a \rangle}{2i \sin \theta} - \varepsilon bD_0(z)a = \\ &= \frac{1}{2i \sin \theta} (1 - \varepsilon bD_0(z)a) (2i \sin \theta + \varepsilon (1 - \varepsilon bD_0(z)a)^{-1} b\langle \cdot, a \rangle). \end{aligned}$$

We are reduced at looking at the kernel of the third factor in the last line. We will show that the following equation admits a solution $\theta = it$ with $t < 0$ close to 0. The singularity we are searching corresponds to solutions of

$$\begin{aligned} 2i \sin \theta + \langle (1 - \varepsilon bD_0(z)a)^{-1} b, a \rangle &= 0 = 2i \sin \theta + \varepsilon \sum_{n=1}^{\infty} \varepsilon^n \langle (bD_0(z)a)^n b, a \rangle \\ &= 2i \sin \theta + \frac{\varepsilon^2}{2} \sum_{\mu, \nu \in \mathbb{Z}} |\mu - \nu| q(\mu) q(\nu) + O(\varepsilon^3) + O(\theta^2). \end{aligned}$$

By the implicit function theorem we obtain a unique solution

$$\theta = \theta(\varepsilon) = i \frac{\varepsilon^2}{4} \sum_{\mu, \nu \in \mathbb{Z}} |\mu - \nu| q(\mu) q(\nu) + O(\varepsilon^2).$$

For $\varepsilon^2 > 0$ we have $\Im\theta(\varepsilon) < 0$. This yields for $\varepsilon^2 > 0$ an eigenvalue $z(\varepsilon) = 2(1 - \cos(\theta(\varepsilon)))$ of H with $z(\varepsilon)$ close to 0. Necessarily $z(\varepsilon) < 0$ by Lemma 5.3 [CT] and by selfadjointness of H . So we obtain $\theta = it$ with $t < 0$ and $t = O(\varepsilon^2)$. This argument proves the existence of an eigenvalue also near 4, since if $(-\Delta + q)u = \lambda u$ then $v(\nu) = (-1)^\nu u(\nu)$ satisfies $(-\Delta - q)v = (4 - \lambda)v$. For ε_0 small, there are no other eigenvalues. We need to show that 0 and 4 are not resonances. We focus on 0. 0 for some ε is a resonance exactly if the Wronskian $W(\varepsilon, \theta)$ is $W(\varepsilon, 0) = 0$ (see §10, we have added the parameter ε). By (10.1) this happens exactly if

$$(1) \quad bR_H(z)a = (1 + \varepsilon bR_{-\Delta}(z)a)^{-1} bR_{-\Delta}(z)a = \\ (2i \sin \theta + \varepsilon(1 - \varepsilon bD_0(z)a)^{-1} b\langle \cdot, a \rangle)^{-1} (1 - \varepsilon bD_0(z)a)^{-1} (2i \sin \theta bR_{-\Delta}(z)a)$$

is singular at $z = 0$. But the first factor in rhs(1) is not singular at $z = \theta = 0$. The second factor is not singular. The third factor has kernel

$$k(\mu, \nu) = b(\mu)a(\nu) - b(\mu)(1 - e^{-i\theta|\mu-\nu|})a(\nu)$$

which is obviously not singular at $\theta = 0$. This means that 0 is not a resonance. The same argument proves also the statement for 4, thanks to the transformation $v(\nu) = (-1)^\nu u(\nu)$. This yields Lemma A.1.

Here we recall that [KKK] observe that if $q(\mu) = A\delta(\mu - \mu_1) + B\delta(\mu - \mu_2)$ with $(A, B) \neq (0, 0)$ then $H = -\Delta + q$ has at least one eigenvalue. In the case of small potentials we can generalize this observation. If $\sum_{\mu \in \mathbb{Z}} q(\mu) \neq 0$ then proceeding as in Lemma A.1 it is easy to show that $H = -\Delta + \varepsilon q$ has an eigenvalue for $\varepsilon \neq 0$ small. For $\sum_{\mu \in \mathbb{Z}} q(\mu) = 0$ we can generally apply Lemma A.1 thanks to the following lemma:

Lemma A.2. *Suppose that $q(\mu) \in \mathcal{S}(\mathbb{Z})$ satisfies $\sum_{\mu \in \mathbb{Z}} q(\mu) = 0$. Then*

$$-\frac{1}{2} \sum_{\mu, \nu \in \mathbb{Z}} |\mu - \nu| q(\mu) q(\nu) = \lim_{z \rightarrow 0^-} \langle R_{-\Delta}(z)q, q \rangle.$$

Notice that for $z < 0$ we have $\langle R_{-\Delta}(z)q, q \rangle > 0$, so the above limit is generically positive. To prove Lemma A.2 use

$$R_{-\Delta}(\mu, \nu, z) = \frac{1}{2i \sin \theta} e^{-i\theta|\mu-\nu|} = \frac{1}{2i \sin \theta} - \frac{\theta|\nu - \mu|}{2 \sin \theta} + O(\theta),$$

We have for $|\cdot| * q(\mu) = \sum_{\nu} |\mu - \nu| q(\nu)$

$$\langle R_{-\Delta}(z)q, q \rangle = \frac{\sum_{\nu} q(\nu)}{2i \sin \theta} - \frac{1}{2} \langle |\cdot| * q, q \rangle + O(\theta) \rightarrow -\frac{1}{2} \langle |\cdot| * q, q \rangle.$$

REFERENCES

- [BV] D.Bambusi, D.Vella, *Quasi periodic breathers in Hamiltonian lattices with symmetries*, Discrete Contin. Dyn. Syst. Ser. B **2** (2002), 389–399.
- [C] S.Cuccagna, *On instability of excited states of the nonlinear Schrödinger equation*, <http://arxiv.org/>.
- [CV] S.Cuccagna, N.Visciglia, *On asymptotic stability of ground states of NLS with a finite bands periodic potential in 1D*, [p://arxiv.org/abs/0809.4775](http://arxiv.org/abs/0809.4775).
- [CT] S.Cuccagna, M.Tarulli, *On asymptotic stability of standing waves of discrete Schrödinger equation in $\mathbb{Z}$* , <http://arxiv.org/abs/0808.2024>.
- [JA] M.Johansson,,S.Aubry, *Existence and stability of quasiperiodic breathers in the discrete nonlinear Schrödinger equation*, Nonlinearity **10** (1997), 1151–1178.
- [KKK] A.Komech, E.Kopylova, M.Kunze, *Dispersive estimates for 1D discrete Schrödinger and Klein Gordon equations*, Appl. Mat. **85** (2006), 1487–1508.
- [KPS] P.G. Kevrekidis, D.E. Pelinovsky, A. Stefanov, *Asymptotic stability of small solitons in the discrete nonlinear Schrödinger equation in one dimension*, <http://arxiv.org/abs/0810.1778>.
- [PS] D.E.Pelinovsky,A. Stefanov, *On the spectral theory and dispersive estimates for a discrete Schrödinger equation in one dimension*, <http://arxiv.org/abs/0804.1963>.
- [RS] M.Reed, B.Simon, *Methods of Mathematical Physics*, Academic Press, San Diego.
- [R] W.Rudin, *Real and Complex Analysis*, third edition, McGraw–Hill, New York.
- [SK] A. Stefanov, P.G.Kevrekidis, *Asymptotic behaviour of small solutions for the discrete nonlinear Schrödinger and Klein–Gordon equations*, Nonlinearity **18** (2005), 1841–1857.

DISMI UNIVERSITY OF MODENA AND REGGIO EMILIA, VIA AMENDOLA 2, PADIGLIONE
MORSELLI, REGGIO EMILIA 42100 ITALY
E-mail address: `cuccagna.scipio@unimore.it`